

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 November 2005 (10.11.2005)

PCT

(10) International Publication Number
WO 2005/105136 A1

(51) International Patent Classification⁷: **A61K 38/21**,
9/00, 9/127, 9/51, A61P 35/00

(21) International Application Number:
PCT/US2005/014626

(22) International Filing Date: 27 April 2005 (27.04.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/565,756 27 April 2004 (27.04.2004) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF SOUTH FLORIDA** [US/US]; 4202 East Fowler Avenue, FAO 126, Tampa, FL 33620-7900 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MOHAPATRA, Shyam, S.** [US/US]; 18510 Country Crest Place, Tampa, FL 33647 (US).

(74) Agents: **LADWIG, Glenn, P.** et al.; Saliwanchik, Lloyd & Saliwanchik, P.O. Box 142950, Gainesville, FL 32614-2950 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **NANOGENE THERAPY FOR CELL PROLIFERATION DISORDERS**

(57) Abstract: The present invention concerns particles comprising a chitin component, such as chitosan or a derivative thereof, associated with a polynucleotide encoding an interferon (IFN) molecule, 2-5' oligoadenylate synthetase (2-5 AS), or a combination thereof. Preferably, the chitin component comprises chitosan or a derivative thereof. The particles of the invention are useful for delivery and expression of the interferon-encoding and/or 2-5 AS-encoding polynucleotide within a host *in vitro* or *in vivo*. The invention further concerns pharmaceutical compositions comprising particles of the invention and a pharmaceutically acceptable carrier, and a method for producing particles of the present invention. The present invention further pertains to a method of inducing apoptosis in a cancer cell, such as a lung cancer cell, by contacting a target cancer cell *in vitro* or *in vivo* with an effective amount of particles of the invention. In one embodiment, a therapeutically effective amount of particles are administered to target cancer cells within a patient *in vivo*, for treatment of cancer, such as lung cancer. The particles and therapeutic methods of the invention provide anti-metastatic and anti-cancer therapeutics for cancer patients, particularly lung cancer patients.



WO 2005/105136 A1

DESCRIPTION

NANOGENE THERAPY FOR CELL PROLIFERATION DISORDERS

5

CROSS-REFERENCE TO RELATED APPLICATION

This application claims benefit of U.S. Provisional Application Serial No. 60/565,756, filed April 27, 2004, which is hereby incorporated by reference herein in its entirety, including any figures, tables, nucleic acid sequences, amino acid sequences, and
10 drawings.

FIELD OF THE INVENTION

This invention pertains to particles including a chitin component, a polynucleotide encoding an interferon molecule, such as IFN-gamma (IFN- γ), or an interferon-inducible
15 molecule such as 2'-5' oligoadenylate synthetase (2-5 AS) or interferon regulatory factor (IRF-1), or a combination of any of the foregoing, and the use of such particles for treatment of cell proliferation disorders, such as lung cancer.

BACKGROUND OF THE INVENTION

20 Lung cancer is one of the leading causes of death worldwide. Despite progress made in our understanding of the multiple risk factors associated with the development of lung cancer, and progress in developing novel approaches, this disease remains difficult to treat effectively. Lung cancer patients often present with locally advanced or disseminated disease. Their long-term survival is poor and such aggressive cancers are
25 difficult to treat because of drug-induced toxicity. Non-viral plasmid DNA (pDNA)-mediated gene therapy, one of several new therapeutic approaches for lung cancer, provides a better alternative that is both safe and effective. Unlike viral vectors, which can induce an immune response with associated immunogenicity and systemic toxicity, a pDNA strategy combined with a chitosan-based nanoparticle (CBN) carrier system
30 provides a unique approach to delivering genes by the mucosal route with limited toxicity and increased transgene expression, especially in target organs such as the lung (disclosed

in Mohapatra *et al.*, international publication WO 03/028759 A1 and Mohapatra *et al.*, U.S. patent publication 2003-0068333-A1, which are each incorporated herein by reference in their entirety).

Lung tumor development and metastasis are complex processes that include transformation, proliferation, resistance to apoptosis, neovascularization, and metastatic spread (Antoniou, K.M. *et al. Chest*, 2003, 123:209-216). A number of gene products have been identified that play critical roles in these processes. Inhibition of metastasis is one of the most important therapeutic strategies in the treatment of lung cancer, since approximately 70% of lung cancer patients die from the metastatic disease even after a complete resection of primary tumor. Metastasis involves the disruption of extracellular matrix (ECM) adhesion, ECM degradation, cell cycle dysregulation, and escape from apoptosis. Thus, protection from metastasis would have to block one or more of these processes.

A complex array of endocrine activities controls cell proliferation and death in the respiratory, gastrointestinal and urinary mucosa, which are major sites of tumor development. Interferons (IFNs) have received wide attention for their anti-cancer effects and are currently used for many cancers. The major oncologic indications of IFNs include melanoma, renal cell carcinoma, AIDS-related Kaposi sarcoma, follicular lymphoma, hairy cell leukemia and chronic myelogenous leukemia (Antoniou, K.M. *et al. Chest*, 2003, 123:209-216). Exogenous recombinant IFNs have a shorter half-life *in vivo*, and systemic administration at moderate to high doses may cause substantial adverse effects (Gutterman, J.U. *PNAS*, 1994, 91:1198-1205; Antoniou, K.M. *et al. Chest*, 2003, 123:209-216).

To overcome the limitations inherent with therapy using cytokines *per se* (cytokine proteins or polypeptides), several investigators have used transient gene expression therapy involving these genes. Separately, IFN- γ and IL-12 have each proven effective both as prophylactics and adjuncts in therapy against diverse human diseases (Mohapatra, S.S. *Science*, 1995, 269(5230):1499; Murray, H.W. *Intensive Care Med*, 1996, 22(Suppl 4):S456-S461). Oromucosal IFN therapy was found to be effective for antiviral and antitumoral activity (Okubo, T. *et al. J Immunol*, 1999, 162:4013-4017). However, mucosal administration of IFN- γ pDNA has not been studied.

The last decade has seen tremendous progress in gene expression technology. Several investigators have utilized a replication-deficient episomal adenovirus as a vehicle for transient gene expression. Adenoviral vectors are very efficient at transducing target cells *in vitro* and *in vivo* and permit transgene expression in a dose-dependent manner (Behera, A.K. *et al. Hum Gene Ther.*, 2002, 13:1697-1709), but they do produce acute inflammation and an immune response to viral vector encoded antigens, which remain the major stumbling blocks to the application of adenovirus-mediated IFN- γ gene transfer for treating human diseases. Previous studies have demonstrated that the mucosal administration of pIFN- γ significantly decreased airway inflammation and airway hyper-responsiveness in a mouse model of grass allergic asthma. Adenoviral-mediated IFN- γ gene transfer effectively reversed established asthma in a BALB/c mouse model (Behera, A.K. *et al. Hum Gene Ther.*, 2002, 13:1697-1709).

It has recently been shown that intranasally delivered pDNA encoding interferon gamma (IFN- γ) can be used as an antiviral treatment against respiratory syncytial virus infection (Mohapatra *et al.*, U.S. Patent No. 6,489,306). Further, IFN- γ is known to induce interferon response factor (IRF-1) and 2'5' oligoadenylate synthetase (2-5 OAS), which also have antiviral properties (Behera, A.K. *et al.*, *JBC*, 2002, 277(28):25601-25608; Mohapatra *et al.*, U.S. patent publication 2004-0009152-A1; Mohapatra *et al.*, international publication WO 03/092618 A2; which are each incorporated herein by reference in their entirety). Also, an IFN- γ producing plasmid encapsulated in a chitin-based nanoparticle, which has been referred to as "CIN", has been shown to possess anti-inflammatory and apoptosis-inducing properties and to attenuate lung inflammation and airway hypereactivity (Kumar *et al. Genet Vacc Ther*, 2003, 1(1):3; Mohapatra, international publication WO 2004/074314 A2; which are each incorporated herein by reference in their entirety).

The present inventors reasoned that intranasally administered nanoparticles capable of *de novo* production of the IFN- γ may provide a novel means of prophylaxis and/or treatment for cancer, such as metastatic lung cancer. Research in the laboratory has identified the pIFN- γ as a potential lung cancer treatment based on its ability to induce significant apoptosis in cultured lung cancer cell lines. Also, CBN complexed pDNA encoding pIFN- γ was found to completely abrogate the development of lung tumors in a nude mouse model of metastatic lung cancer.

Non-viral mediated gene expression using plasmid DNAs (pDNAs) has a number of advantages, including ease of preparation and use, stability, and room temperature storage (Hellerman, G.R. and Mohapatra, S.S. *Gen Vacc & Ther*, 2003, 1:1). They do not replicate in mammalian cells and do not integrate into host genomes, yet they can persist in host cells and express the cloned gene for a period of weeks to months. One problem associated with the pDNA approach is inefficient gene transfer *in vivo*, especially in slow and non-dividing cells such as epithelial cells (Mohapatra, S.S. *Pediatr. Infect. Dis. J.*, 2003, 22(2 Suppl):S100-S103). CBNs protect pDNA from nuclease degradation and facilitate its entry into target cells. CBNs are prepared from chitosan, a biocompatible cationic polysaccharide from chitin extracted from crustacean shells, and have shown excellent potential for gene (Mao, H-Q. *et al. J. Controlled Release*, 2001, 70(3):399-421, which is incorporated herein by reference in its entirety) and controlled drug delivery. Chitosan is non-toxic, resistant to biodegradation, non-hemolytic, stimulates the immune system, is an anticoagulant, and has wound-healing and antimicrobial properties. Chitosan also increases transcellular and paracellular transport across the mucosal epithelium, thereby facilitating mucosal gene delivery. Another advantage of the use of CBNs for gene transport is their ability to target specific cells. Reduction of nonspecific interactions by shielding of net positive surface charges also improves targeting of CBNs.

BRIEF SUMMARY OF THE INVENTION

In one aspect, the present invention concerns particles comprising a chitin component, which is associated with a polynucleotide encoding an interferon (IFN) or an IFN-inducible protein, such as 2'-5' oligoadenylate synthetase (2-5 AS) or interferon regulatory factor (IRF-1). Preferably, the chitin component comprises chitosan or a derivative thereof. Optionally, the particles of the invention further comprise a lipid component and are referred to herein interchangeably as "chliposomes", "chlipids", "chitosan-lipid nanoparticles" or "CLNs". The particles of the invention are useful for delivery and expression of the interferon-encoding and/or IFN-inducible molecule-encoding polynucleotide within a host *in vitro* or *in vivo*. The invention further concerns a method for producing particles of the present invention.

In some embodiments, the particles of the invention comprise a polynucleotide encoding an interferon selected from the group consisting of alpha-interferon, beta-

interferon, gamma-interferon, omega-interferon, and lambda-interferon. In some embodiments, the particles of the invention comprise a polynucleotide encoding 2-5 AS or at least one catalytically active fragment thereof selected from the group consisting of the p40, p69, and p100 subunit. Such 2-5 AS subunits may be one or more splice
5 variants, such as the 42kDa, 46kDa, 69kDa, and/or 71kDa variant. In some embodiments, the particles of the invention comprise a polynucleotide encoding IRF-1, or a biologically active fragment or homolog thereof.

In another aspect, the present invention concerns a pharmaceutical composition comprising particles comprising a chitin component and a polynucleotide encoding an
10 interferon (IFN) molecule, an IFN-inducible molecule, or a combination thereof; and a pharmaceutically acceptable carrier. Optionally, the particles of the invention further comprise a lipid component. In one embodiment, the pharmaceutical composition is formulated for delivery through a mucosal route, such as the lungs.

In another aspect, the present invention concerns a method of treating a cell
15 proliferation disorder by administering a therapeutically effective amount of particles to a patient in need thereof. Accordingly, a method of reducing cellular growth by administering a therapeutically effective amount of particles of the invention is contemplated, in order to reduce (partially or completely inhibit, prevent, or slow) uncontrolled cell growth. In one embodiment, an effective amount of particles are
20 administered to a patient for treatment of cancer, such as lung cancer.

In another aspect, the present invention concerns a method of inducing apoptosis in a cancer cell, such as a lung cancer cell, by contacting a target cancer cell *in vitro* or *in vivo* with an effective amount of particles of the invention. In one embodiment, a therapeutically effective amount of particles are administered to target cancer cells within
25 a patient *in vivo*, for treatment of cancer, such as lung cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be
30 provided by the Office upon request and payment of the necessary fee.

Figure 1 shows an analysis of apoptosis using fluorescence microscopy in cells transfected with pIFN- γ . The micrographs show that IFN-gamma treatment of HEp-2 cells induces apoptosis.

Figure 2 shows an immunoblot demonstrating the detection of p27kip expression and PARP cleavage in IFN-gamma treated HEp-2 cells with or without RSV infection.

Figure 3 shows immunocytochemistry following pIFN-gamma treatment. BALB/c nude mice were injected with A549 cells (5×10^6 cells/mouse) intravenously (i.v.) and one group treated with pIFN-gamma and another group with pVAX as control.

Figures 4A-4D show derivation and characterization of NG-042 nanoparticles. Figure 4A shows synthesis and characterization of nanochitosan particles produced by proprietary method. The products were separated by capillary gel electrophoresis. The plot shows the separation of 4 low molecular weight components. The nanogene particles were then subject to analysis of size and zeta potential using a NiComp381 Zetasizer. Results are shown in Figure 4B. The intensity weight distribution of NG042 particles showing their size of 155 nm, zeta potential=20.42. Atomic Force Microscopic analysis of Nanogene-042 particles showing oligomeric structure complexed with DNA (red arrows; upper line) is shown in Figure 4C. Figure 4D shows that lyophilized and resuspended NG042 particles retain functionality at ambient temperatures of 23° to 55° C. Nanogene complexes of pGL3 (firefly luciferase, Promega) was lyophilized, reconstituted with water and treated for 24 hours at RT (23° C), 42° C, 55° C and -20° C. A549 cells were plated and transfected with the above complexes. Uptake and expression of DNA was allowed to occur for 24 hours. Luciferase activity was determined by using Promega's Dual Assay kit. Readings were normalized to relative luminiscence units (RLU) per mg protein.

Figures 5A-5C show characterization of NG044 particles. Figure 5A shows that expression of nanoparticle-encapsulated EGFP gene continues *in vivo* until day 10. NG044 particles were complexed with DNA (5:1) encoding green fluorescent protein and administered intranasally to groups of mice (n=3). Mice were sacrificed on the indicated days and broncho-alveolar lavage cells were examined by fluorescent microscopy. Figure 5B demonstrates the thermogelling property of NG044. NG044 forms a gel upon reacting with 2-glycerol phosphate, while NG042, another depolymerized chitosan, does not. To test the controlled release of gene expression, NG044 hydrogel was prepared

using pEGFP plasmid DNA and PVP/glutaraldehyde for gel formation. The hydrogel was freeze-dried and the powder was resuspended in water (NG044 hydrogel) and given intranasally to groups of mice (n=4). Another group received NG044 with pEGFP without gelling (Control). Gene expression in the mouse lung was measured by EGFP expression in BAL cells 10 and 20 days after administration. Results are shown in Figure 5C. The results at day 10 were similar (not shown) for control and hydrogel, whereas after 20 days mice given hydrogel continued EGFP show expression and no expression was detected in control mice.

10

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO: 1 is a nucleotide coding sequence (CDS) for the human 40kDa splice variant of the 40/46kDa subunit ("p40 subunit") of 2'-5' oligoadenylate synthetase (National Center for Biotechnology Information (NCBI) Accession Number NM_016816).

15

SEQ ID NO: 2 is an amino acid sequence of the human 40kDa splice variant of the 40/46kDa subunit ("p40 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_016816).

20

SEQ ID NO: 3 is a nucleotide coding sequence (CDS) for the human 46kDa splice variant of the 40/46kDa subunit ("p40 subunit") of 2'-5' oligoadenylate synthetase (National Center for Biotechnology Information (NCBI) Accession Number NM_016816).

25

SEQ ID NO: 4 is an amino acid sequence of the human 46kDa splice variant of the 40/46kDa subunit ("p40 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_016816).

SEQ ID NO: 5 is a nucleotide coding sequence (CDS) for the human 69 kDa splice variant of the 69/71kDa subunit ("p69 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_002535).

30

SEQ ID NO: 6 is an amino acid sequence of the human 69 kDa splice variant of the 69/71kDa subunit ("p69 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_002535).

SEQ ID NO: 7 is a nucleotide coding sequence (CDS) for the human 71 kDa splice variant of the 69/71kDa subunit ("p69 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_002535).

SEQ ID NO: 8 is an amino acid sequence of the human 71kDa splice variant of the 69/71kDa subunit ("p69 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number NM_002535).

SEQ ID NO: 9 is a nucleotide coding sequence (CDS) for the human 100kDa subunit ("p100 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number AF063613).

SEQ ID NO: 10 is an amino acid sequence of the human 100kDa subunit ("p100 subunit") of 2'-5' oligoadenylate synthetase (NCBI Accession Number AF063613).

SEQ ID NO: 11 is a nucleotide coding sequence (CDS) for the mouse homolog of the 2'-5' oligoadenylate synthetase 40kDa splice variant (p40 subunit) (NCBI Accession Number M33863).

SEQ ID NO: 12 is the amino acid sequence for the mouse homolog of the 2'-5' oligoadenylate synthetase 40kDa splice variant (p40 subunit) (NCBI Accession Number M33863).

SEQ ID NO: 13 is the human 2'-5' oligoadenylate synthetase 40/46kDa (p40 subunit) gene (NCBI Accession Number NM_016816).

SEQ ID NO: 14 is the human 2'-5' oligoadenylate synthetase 69/71kDa (p69 subunit) gene (NCBI Accession Number NM_002535).

SEQ ID NO: 15 is the human 2'-5' oligoadenylate synthetase 100kDa (p100 subunit) gene (NCBI Accession Number AF063613).

SEQ ID NO: 16 is the mouse homolog of the 2'-5' oligoadenylate synthetase 40kDa (p40 subunit) gene (NCBI Accession Number M33863).

SEQ ID NO: 17 is the nucleotide coding sequence (CDS) for human IFN- γ (NCBI Accession No: NM_000639).

SEQ ID NO: 18 is the amino acid sequence for human IFN- γ (NCBI Accession No: NM_000639).

SEQ ID NO: 19 is the nucleotide coding sequence (CDS) for human interferon-beta (NCBI Accession No.: M25460).

SEQ ID NO:20 is the nucleotide coding sequence (CDS) for human interferon-beta-1 (NCBI Accession No.: M28622).

SEQ ID NO:21 is the nucleotide coding sequence (CDS) for a human interferon (NCBI Accession No.: L25664).

5 **SEQ ID NO:22** is the nucleotide coding sequence (CDS) for human interferon-alpha (NCBI Accession No.: M54886 and M38682).

SEQ ID NO:23 is the nucleotide coding sequence (CDS) for human interferon-alpha-J1 (NCBI Accession No.: M34913).

10 **SEQ ID NO:24** is the nucleotide coding sequence (CDS) for human interferon-omega-1 (NCBI Accession No.: X58822).

SEQ ID NO:25 is the nucleotide coding sequence (CDS) for human interleukin 28A (interferon, lambda 2; IL-28A) (NCBI Accession No.: NM_172138).

SEQ ID NO:26 is the nucleotide coding sequence (CDS) for human interleukin 28B (interferon lambda 3; IL-28B) (NCBI Accession No.: AY336714).

15 **SEQ ID NO:27** is the nucleotide coding sequence (CDS) for human interleukin 28C (interferon lambda 4; IL-28C) (NCBI Accession No.: AY336717).

SEQ ID NO:28 is the nucleotide coding sequence (CDS) for human interleukin 29 (interferon lambda 1; IL-29) (NCBI Accession No.: NM_172140).

20 **SEQ ID NO:29** is the nucleotide coding sequence (CDS) for a human interferon-like peptide (NCBI Accession No.: EE00870).

SEQ ID NO:30 is the nucleotide coding sequence (CDS) for a human interferon-like peptide (NCBI Accession No.: EE00871).

SEQ ID NO:31 is the nucleotide coding sequence (CDS) for a human interferon-regulatory factor 1 (IRF-1) (NCBI Accession No.: 002198).

25

DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns particles comprising a chitin component, such as chitosan or a derivative thereof, associated with a polynucleotide encoding an interferon (IFN) molecule or an IFN-inducible molecule, or a combination thereof. Preferably, the particles further comprise a control sequence operably-linked to the polynucleotide, which is capable of causing expression of the polynucleotide within a host *in vitro* or *in vivo*.

30

In certain embodiments, the interferon molecule encoded by the polynucleotide is Type I or Type II interferon, including those commonly designated as alpha-interferon, beta-interferon, gamma-interferon, and omega-interferon (also designated α -interferon, β -interferon, γ -interferon, and ω -interferon), and combinations thereof, including the
5 consensus sequence for alpha-interferon. In some embodiments, the alpha-interferon is α_1 or α_2 -interferon. In some embodiments, the interferon is interferon α -2b. Other interferons include interferon α -2 β , a fusion interferon α -2 α -1, interferon α -2e, human $\alpha 1$ or $\alpha 2$ interferon.

In some embodiments, the interferon is a hybrid interferon. The construction of
10 hybrid polynucleotides encoding combinations of different interferon subtypes (such as α and ϵ ; α and β , and α and F) is disclosed in U.S. Patent Nos. 4,414,150; 4,456,748; and 4,678,751, each of which are incorporated herein by reference in their entirety. U.S. Patent Nos. 4,695,623; 4,897,471; and 5,831,062, which are incorporated herein by
15 reference in their entirety, disclose novel human leukocyte interferon polypeptides having amino acid sequences that include common or predominant amino acids found at each position among naturally-occurring alpha interferon subtype polypeptides and are referred to as consensus human leukocyte interferon. In one embodiment of the invention, the hybrid interferon is interferon $\alpha 2\alpha 1$.

In one embodiment, the interferon is an interferon- α . Recombinant interferon
20 alphas, for instance, have been cloned and expressed in *E. coli* by several groups (*e.g.*, Weissmann *et al.*, *Science*, 1980, 209:1343-1349; Sreuli *et al.*, *Science*, 1980, 209:1343-1347; Goeddel *et al.*, *Nature*, 1981, 290:20-26; Henco *et al.*, *J. Mol. Biol.*, 1985, 185:227-260, each of which are incorporated herein by reference in their entirety). In some embodiments, the interferon is a human interferon alpha. In some embodiments, the
25 interferon alpha is interferon alpha 2a or 2b.

The term "interferon" as used herein is intended to include all classes and subclasses of interferon, and deletion, insertion, or substitution variants, as well as "interferon-like" molecules such as interleukin 15 (IL-15), interleukin 28A (interferon
30 lambda2; IL-28A), interleukin 28B (IL-28B), interleukin 28C (IL-28C), interleukin 29 (interferon lambda1; IL-29), and synthetic interferon-like peptides (*e.g.*, NCBI accession nos. E00871 and E00870). In one embodiment, the interferon-encoding polynucleotide, or its polypeptide product, is the interferon-alpha-encoding polynucleotide or its

polypeptide product. In some embodiments, the interferon-encoding polynucleotide of the particle, or its polypeptide, is the human nucleotide or amino acid sequence. The human interferon alphas, for example, are a family of proteins including at least 24 subspecies (Zoon, K. C., *Interferon*, 1987, 9:1, Gresser, I., ed., Academic Press, NY).

5 The interferon alphas were originally described as agents capable of inducing an antiviral state in cells but are now known as pleiotropic lymphokines affecting many functions of the immune system (Openakker *et al.*, *Experimentia*, 1989, 45:513). In some embodiments, the interferon alpha is interferon alpha 2a or 2b (see, for example, WO 91/18927, which is incorporated by reference herein in its entirety), although any
10 interferon alpha may be used. Nucleotide sequences encoding the exemplified interferons interferon-gamma; interferon-beta; interferon-beta-1; interferon; interferon-alpha; interferon-alpha-J1; interferon omega-1; interleukin 28A; interleukin 28B; interleukin 28C, interleukin 29; and interferon-like peptides are listed as SEQ ID NOs: 17 and 19-30. Particles of the invention may contain one or more of these polynucleotides or degenerate
15 sequences encoding the same polypeptides, for example

The interferon-encoding polynucleotide may encode gamma-interferon (IFN- γ), among others. IFN- γ is a 14-18 kDalton 143 amino acid glycosylated protein that is a potent multifunctional cytokine. As used herein, "interferon-gamma", "IFN-gamma", "interferon- γ ", and "IFN- γ " refer to IFN- γ protein, biologically active fragments of IFN- γ ,
20 and biologically active homologs of "interferon-gamma" and "IFN- γ ", such as mammalian homologs. These terms include IFN- γ -like molecules. An "IFN- γ -like molecule" refers to polypeptides exhibiting IFN- γ -like activity when the polynucleotide encoding the polypeptide is expressed, as can be determined *in vitro* or *in vivo*. For purposes of the subject invention, IFN- γ -like activity refers to those polypeptides having
25 one or more of the functions of the native IFN- γ cytokine disclosed herein (such as induction of apoptosis). Fragments and homologs of IFN- γ retaining one or more of the functions of the native IFN- γ cytokine, such as those disclosed herein, is included within the meaning of the term "IFN- γ ". In addition, the term includes a nucleotide sequence which through the degeneracy of the genetic code encodes a similar peptide gene product
30 as IFN- γ and has the IFN- γ activity described herein. For example, a homolog of "interferon-gamma" and "IFN- γ " includes a nucleotide sequence which contains a "silent" codon substitution (*e.g.*, substitution of one codon encoding an amino acid for

another codon encoding the same amino acid) or an amino acid sequence which contains a “silent” amino acid substitution (*e.g.*, substitution of one acidic amino acid for another acidic amino acid).

An exemplified nucleotide sequence encodes human IFN- γ (Accession No:
5 NM_000639, NCBI database, which is hereby incorporated by reference in its entirety):

```
1 tgaagatcag ctattagaag agaaagatca gttaagtctt ttggacctga tcagcttgat
61 acaagaacta ctgatttcaa cttctttggc ttaattctct cggaacgat gaaatataca
121 agttatatct tggttttca gctctgcac gtttgggtt ctctggctg ttactgccag
10 181 gaccatattg taaaagaagc agaaaacctt aagaaatatt ttaatgcagg tcattcagat
241 gtatcgata atggaactct ttcttaggc atttgaaga attggaaga ggagagtgc
301 agaaaaataa tgcagagcca aattgtctcc ttctactca aacttttaa aaactttaa
361 gatgaccaga gcatccaaa gagtgtggag accatcaagg aagacatgaa tgtcaagttt
421 ttcaatagca acaaaaagaa acgagatgac ttcgaaaagc tgactaatta ttcggtaact
15 481 gacttgaatg tccaacgcaa agcaatacat gaactcatcc aagtgatggc tgaactgtcg
541 ccagcagcta aaacagggaa gcgaaaagg agtcagatgc tgttcaagg tcgaagagca
601 tccagtaat ggtgtcctg cctgcaatat ttgaattta aatctaaac tattattaa
661 tatttaacat tatttatatg gggaatatat tttagactc atcaatcaa taagtattta
721 taatagcaac tttgtgtaa tgaaaatgaa tatctattaa tatatgtatt attataatt
20 781 cctatacct gtgactgtct cacttaatcc ttgttttct gactaattag gcaaggctat
841 gtgattacaa ggctttatct caggggcca ctaggcagcc aacctagca agatcccatg
901 ggttgtgtgt ttattcact tgatgataca atgaacactt ataagtgaag tgatactatc
961 cagttactgc cggttgaaa atatgcctgc aatctgagcc agtgcttaa tggcatgtca
1021 gacagaactt gaatgtgtca ggtgacctg atgaaacat agcatctcag gagatttcat
25 1081 gcctggtgct tccaaatatt gtgacaact gtgactgtac ccaaatggaa agtaactcat
1141 ttgttaaata tatcaatac taatatatat gaataaagtg taagtcaca act (SEQ ID NO: 17)
```

```
MKYTSYILAFQLCIVLGS LGSCYQDPYVKEAENLKKYFNAGHSDVADNGTLFLG
ILKNWKEESDRKIMQSQIVSFYFKLFKNFKDDQSIQKSVETIKEDMNVKFFNSNK
30 KKRDDFEKLTNYSVTDLNVQRKAIHELIQVMAELSPA AKTGKRKRSQMLFQ
GRRASQ (SEQ ID NO: 18)
```

U.S. Patent Nos. 5,770,191 and 6,120,762, which are incorporated herein by reference in their entirety, describe several C-terminal fragments of IFN-gamma that may be encoded by the polynucleotide(s) carried by the particles of the invention. Other interferons that may be encoded by polynucleotides within the particles of the invention are described in U.S. patent publications 2005-0054052-A1; 2005-0054053-A1; 2005-0025742-A1; and 2005-0084478-A1; which are incorporated herein by reference in their entirety.

Interferon regulatory factor-1 (IRF-1) is an interferon-inducible molecule (*e.g.*, an interferon-stimulated gene product) (Pizzoferrato, F. *et al.*, *Cancer Res.*, 2004, 64(22):8381-8388; Pack, S.Y. *et al.*, *Eur. J. Biochem.*, 2004, 271(21):4222-4228). The nucleotide sequence encoding human IRF-1 is provided herein as SEQ ID NO: 31. Particles of the invention may contain this polynucleotide, degenerate sequences encoding the same polypeptide, or biologically active fragments thereof, for example.

2'5' oligoadenylate synthetase (2-5 AS) is an interferon-inducible molecule. In some embodiments, the particles of the invention comprise a polynucleotide encoding 2-5 AS or at least one catalytically active fragment thereof selected from the group consisting of the p40, p69, and p100 subunit. Such 2-5 AS subunits may be one or more splice variants, such as the 42kDa, 46kDa, 69kDa, and/or 71kDa variant. For example, the particles can comprise one or more nucleotide sequences encoding polypeptides comprising one more amino acid sequences set forth herein as SEQ ID NOs: 2, 4, 5, 6, 8, 10, 12, 13, 14, 15, or 16, or catalytically active fragments of these amino acids. In some embodiments, the particles comprise one or more nucleotide sequences selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 14, 15, 16, and 18, or a catalytically active fragment thereof.

As indicated above, the particle used in the compositions and methods of the invention comprise polynucleotides encoding an interferon, an interferon-inducible molecule, or both. For example, the particle can contain a polynucleotide encoding an interferon and 2-5 AS; encoding an interferon and IRF-1; encoding 2-5 AS and IRF-1; or encoding an interferon, 2-5 AS, and IRF-1. Combinations of an interferon and interferon-inducible molecules can be encoded by polynucleotides within a single particle or multiple different particles.

The nucleotide sequences encoding interferon and/or an interferon-inducible molecule used in the subject invention include “homologous” or “modified” nucleotide sequences. Modified nucleic acid sequences will be understood to mean any nucleotide sequence obtained by mutagenesis according to techniques well known to persons skilled in the art, and exhibiting modifications in relation to the normal sequences. For example, mutations in the regulatory and/or promoter sequences for the expression of a polypeptide that result in a modification of the level of expression of a polypeptide according to the invention provide for a “modified nucleotide sequence”. Likewise, substitutions, deletions, or additions of nucleic acids to the polynucleotides of the invention provide for “homologous” or “modified” nucleotide sequences. In various embodiments, “homologous” or “modified” nucleic acid sequences have substantially the same biological or serological activity as the native (naturally occurring) interferon and/or interferon-inducible polypeptide. A “homologous” or “modified” nucleotide sequence will also be understood to mean a splice variant of the polynucleotides of the instant invention or any nucleotide sequence encoding a “modified polypeptide” as defined below.

A homologous nucleotide sequence, for the purposes of the present invention, encompasses a nucleotide sequence having a percentage identity with the bases of the nucleotide sequences of between at least (or at least about) 20.00% to 99.99% (inclusive). The aforementioned range of percent identity is to be taken as including, and providing written description and support for, any fractional percentage, in intervals of 0.01%, between 20.00% and 99.99%. These percentages are purely statistical and differences between two nucleic acid sequences can be distributed randomly and over the entire sequence length.

In various embodiments, homologous sequences exhibiting a percentage identity with the bases of the nucleotide sequences of the present invention can have 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identity with the polynucleotide sequences of the instant invention. Homologous nucleic acid sequences and amino acid sequences include mammalian homologs of the human interferon and/or interferon-inducible molecule

nucleic acid sequences and amino acid sequences, including homologs of biologically active fragments, such as biologically active subunits.

Both protein and nucleic acid sequence homologies may be evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman *Proc. Natl. Acad. Sci. USA*, 1988, 85(8):2444-2448; Altschul *et al. J. Mol. Biol.*, 1990, 215(3):403-410; Thompson *et al. Nucleic Acids Res.*, 1994, 22(2):4673-4680; Higgins *et al. Methods Enzymol.*, 1996, 266:383-402; Altschul *et al. J. Mol. Biol.*, 1990, 215(3):403-410; Altschul *et al. Nature Genetics*, 1993, 3:266-272).

Nucleotide sequences encoding polypeptides with enhanced interferon activity or interferon-inducible molecule activity (such as 2-5 AS catalytic activity and/or IRF-1 activity) can be obtained by “gene shuffling” (also referred to as “directed evolution”, and “directed mutagenesis”), and used in the compositions and methods of the present invention. Gene shuffling is a process of randomly recombining different sequences of functional genes (recombining favorable mutations in a random fashion) (U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; and 5,837,458). Thus, protein engineering can be accomplished by gene shuffling, random complex permutation sampling, or by rational design based on three-dimensional structure and classical protein chemistry (Cramer *et al., Nature*, 391:288-291, 1998; and Wulff *et al., The Plant Cell*, 13:255-272, 2001)

Identity and similarity of related nucleic acid molecules and polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; York (1988); Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; York (1993); Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Jersey (1994); Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; York (1991); and Carillo *et al., SIAM J. Applied Math.*, 1988, 48:1073.

The particles, methods, and compositions of the present invention can utilize biologically active fragments of nucleic acid sequences encoding interferon and/or

interferon-inducible molecules. Representative fragments of the polynucleotide sequences according to the invention will be understood to mean any polynucleotide fragment having at least 8 or 9 consecutive nucleotides, preferably at least 12 consecutive nucleotides, and still more preferably at least 15 or at least 20 consecutive nucleotides of the sequence from which it is derived. The upper limit for such fragments is the total number of nucleotides found in the full-length sequence (or, in certain embodiments, of the full length open reading frame (ORF) identified herein).

In other embodiments, fragments can comprise consecutive nucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, and up to one nucleotide less than the full-length interferon and/or 2-5 AS coding sequences (or, in some embodiments, up to the full length of nucleotides in the open reading frame (ORF)).

In some embodiments, fragments comprise biologically active subunits (such as the p40 subunit of 2-5 AS (e.g., 40kDa, 42kDa, 46kDa, or other splice variant), p69 subunit of 2-5 AS (e.g., 69 kDa, 71 kDa, or other splice variant), p100 subunit of 2-5 AS, or combinations thereof).

It is also well known in the art that restriction enzymes can be used to obtain biologically active fragments of nucleic acid sequences, such as those encoding interferon and/or interferon-inducible molecules. For example, *Bal31* exonuclease can be conveniently used for time-controlled limited digestion of DNA (commonly referred to as "erase-a-base" procedures). See, for example, Maniatis *et al.* (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York; Wei *et al.* [1983] *J. Biol. Chem.* 258:13006-13512.

Optionally, each particle of the invention further comprises a lipid component that is complexed with the chitin and polynucleotide components of the particle. Since efficient gene expression *in vivo* requires both complex formation for cell uptake and prevention of nucleotide degradation and complex dissociation for transcription by RNA polymerase, a combination of a chitin component and lipid component (such as chitosan

and liposomes, respectively) may lead to increased gene delivery and expression *in vivo*. Therefore, this embodiment of the particle combines these two different carrier systems (also referred to herein interchangeably as “chliposomes”, “chlipids”, “chitosan-lipid nanoparticles” or “CLNs”) to significantly increase polynucleotide transfection and expression. Preferably, the components of the chlipid are oriented such that the polynucleotide is surrounded by a lipid monolayer, with polynucleotide-lipid inverted cylindrical micelles arranged in a hexagonal lattice. Methods for producing CLNs containing polynucleotides encoding interferon-gamma are described in Mohapatra *et al.* international publication WO 2004/074314 A2, which is hereby incorporated herein in its entirety.

The present invention further includes a method for producing the particles of the invention by mixing (*e.g.*, complexing) a polynucleotide and a chitin component, such as chitosan or a chitosan derivative, to form a particle comprising a binary complex of the polynucleotide and the chitin component. Optionally, the method further comprises mixing (complexing) a lipid with the polynucleotide and chitin component to form a particle (CLN) comprising a multiplex of the polynucleotide, the chitin component, and the lipid. Typically, the particles of the present invention range in size from the nanometer range (*e.g.*, less than one micrometer; nanoparticles) to the micrometer size range (*e.g.*, about one micrometer or larger). Methods for producing chitosan-based DNA particles are described in Mohapatra, S.S. *Pediatr. Infect. Dis. J.*, 2003, 22(2 suppl.):S100-S103; Kumar, M. *et al.*, *Hum. Gene Ther.*, 2002, 13(12):1415-1425; Kumar *et al.*, *Genetic Vaccines and Therapy*, 2003, 1:3; and Mohapatra *et al.*, international publication no. WO 2004/074314 A2; each of which are incorporated herein by reference in their entirety.

The type of reaction vessel or substrate utilized for producing the particles of the present invention, or the size of the vessel or substrate, is not critical. Any vessel or substrate capable of holding or supporting the reactants so as to allow the reaction to take place can be used. It should be understood that, unless expressly indicated to the contrary, the terms “adding”, “contacting”, “mixing”, “reacting”, “combining” and grammatical variations thereof, are used interchangeably to refer to the mixture of reactants of the method of the present invention (such as plasmid DNA or a non-

polynucleotide agent such as chitosan or a chitosan derivative, lipid, and so forth), and the reciprocal mixture of those reactants, one with the other (*i.e.*, vice-versa), in any order.

It will be readily apparent to those of ordinary skill in the art that a number of general parameters can influence the efficiency of transfection or polynucleotide delivery.

5 These include, for example, the concentration of polynucleotide to be delivered, the concentration of the chitin component (such as chitosan or a chitosan derivative), and the concentration of lipid (for chlipids of the present invention). For *in vitro* delivery, the number of cells transfected, the medium employed for delivery, the length of time the cells are incubated with the particles of the invention, and the relative amount of particles
10 can influence delivery efficiency. For example, a 1:5 ratio of polynucleotide to lipid, 1:5 ratio of polynucleotide to chitosan, and 20% serum is suitable. These parameters can be optimized for particular cell types and conditions. Such optimization can be routinely conducted by one of ordinary skill in the art employing the guidance provided herein and knowledge generally available to those skilled in the art. It will also be apparent to those
15 of ordinary skill in the art that alternative methods, reagents, procedures and techniques other than those specifically detailed herein can be employed or readily adapted to produce the particles and compositions of the invention. Such alternative methods, reagents, procedures and techniques are within the spirit and scope of this invention.

In accordance with the present invention, the polynucleotides carried by the
20 particles are conjugated with a chitin component, such as chitosan or chitosan derivatives. For example, DNA chitosan nanospheres can be generated, as described by Roy, K. *et al.* (*Nat Med*, 1999, 5:387). Chitosan allows increased bioavailability of the nucleic acid sequences because of protection from degradation by serum nucleases in the matrix and thus has great potential as a mucosal gene delivery system. Chitosan also has many
25 beneficial effects, including anticoagulant activity, wound-healing properties, and immunostimulatory activity, and is capable of modulating immunity of the mucosa and bronchus-associated lymphoid tissue.

The term "chitosan", as used herein, will be understood by those skilled in the art to include all derivatives of chitin, or poly-N-acetyl-D-glucosamine (including all
30 polyglucosamine and oligomers of glucosamine materials of different molecular weights), in which the greater proportion of the N-acetyl groups have been removed through hydrolysis. Generally, chitosans are a family of cationic, binary hetero-polysaccharides

composed of (1→4)-linked 2-acetamido-2-deoxy-β-D-glucose (GlcNAc, A-unit) and 2-amino-2-deoxy-β-D-glucose, (GlcN; D-unit) (Varum K.M. *et al.*, *Carbohydr. Res.*, 1991, 217:19-27; Sannan T. *et al.*, *Macromol. Chem.*, 1776, 177:3589-3600). Preferably, the chitosan has a positive charge. Chitosan, chitosan derivatives or salts (*e.g.*, nitrate, phosphate, sulphate, hydrochloride, glutamate, lactate or acetate salts) of chitosan may be used and are included within the meaning of the term “chitosan”. As used herein, the term “chitosan derivatives” are intended to include ester, ether or other derivatives formed by bonding of acyl and/or alkyl groups with OH groups, but not the NH₂ groups, of chitosan. Examples are O-alkyl ethers of chitosan and O-acyl esters of chitosan. Modified chitosans, particularly those conjugated to polyethylene glycol, are included in this definition. Low and medium viscosity chitosans (for example CL113, G210 and CL110) may be obtained from various sources, including PRONOVA Biopolymer, Ltd. (UK); SEIGAGAKU America Inc. (Maryland, USA); MERON (India) Pvt, Ltd. (India); VANSOON Ltd. (Virginia, USA); and AMS Biotechnology Ltd. (UK). Suitable derivatives include those which are disclosed in Roberts, Chitin Chemistry, MacMillan Press Ltd., London (1992). Optimization of structural variables such as the charge density and molecular weight of the chitosan for efficiency of polynucleotide delivery and expression is contemplated and encompassed by the present invention.

The chitosan (or chitosan derivative or salt) used preferably has a molecular weight of 4,000 Dalton or more, preferably in the range 25,000 to 2,000,000 Dalton, and most preferably about 50,000 to 300,000 Dalton. Chitosans of different low molecular weights can be prepared by enzymatic degradation of chitosan using chitosanase or by the addition of nitrous acid. Both procedures are well known to those skilled in the art and are described in various publications (Li *et al.*, *Plant Physiol. Biochem.*, 1995, 33: 599-603; Allan and Peyron, *Carbohydrate Research*, 1995, 277:257-272; Damard and Cartier, *Int. J. Biol. Macromol.*, 1989, 11: 297-302). Preferably, the chitosan is water-soluble and may be produced from chitin by deacetylation to a degree of greater than 40%, preferably between 50% and 98%, and more preferably between 70% and 90%.

The lipid component utilized for the particles, compositions, and methods of the present invention is preferably a phospholipid or cationic lipid. Cationic lipids are amphipathic molecules, containing hydrophobic moieties such as cholesterol or alkyl side chains and a cationic group, such as an amine. Phospholipids are amphipathic molecules

containing a phosphate group and fatty acid side chains. Phospholipids can have an overall negative charge, positive charge, or neutral charge, depending on various substituents present on the side chains. Typical phospholipid hydrophilic groups include phosphatidyl choline, phosphatidylglycerol, and phosphatidyl ethanolamine moieties.

5 Typical hydrophobic groups include a variety of saturated and unsaturated fatty acid moieties. The lipids used in the present invention include cationic lipids that form a complex with the genetic material (*e.g.*, polynucleotide), which is generally polyanionic, and the chitosan or chitosan derivative. The lipid may also bind to polyanionic proteoglycans present on the surface of cells. The cationic lipids can be phospholipids or
10 lipids without phosphate groups.

A variety of suitable cationic lipids are known in the art, such as those disclosed in International Publication No. WO 95/02698, the disclosure of which is herein incorporated by reference in its entirety. Exemplified structures of cationic lipids useful in the particles of the present invention are provided in Table 1 of International
15 Publication No. WO 95/02698. Generally, any cationic lipid, either monovalent or polyvalent, can be used in the particles, compositions and methods of the present invention. Polyvalent cationic lipids are generally preferred. Cationic lipids include saturated and unsaturated allyl and alicyclic ethers and esters of amines, amides or derivatives thereof. Straight-chain and branched alkyl and alkene groups of cationic
20 lipids can contain from 1 to about 25 carbon atoms. Preferred straight-chain or branched alkyl or alkene groups have six or more carbon atoms. Alicyclic groups can contain from about 6 to 30 carbon atoms. Preferred alicyclic groups include cholesterol and other steroid groups. Cationic lipids can be prepared with a variety of counterions (anions) including among others: chloride, bromide, iodide, fluoride, acetate, trifluoroacetate,
25 sulfate, nitrite, and nitrate.

Transfection efficiency can be increased by using a lysophosphatide in particle formation. Preferred lysophosphatides include lysophosphatidylcholines such as 1-oleoyllysophosphatidylcholine and lysophosphatidylethanolamines. Well known lysophosphatides which may be used include DOTMA (dioleoyloxypropyl trimethylammonium chloride/DOPE (*i.e.*, LIPOFECTIN, GIBCO/BRL, Gaithersburg,
30 Md.), DOSPA, (dioleoyloxy sperminecarboxamidoethyl dimethylpropanaminium trifluoroacetate)/DOPE (*i.e.*, LIPOFECTAMINE), LIPOFECTAMINE 2000, and DOGS

(dioctadecylamidospemine) (*i.e.*, TRANSFECTAM), and are all commercially available. Additional suitable cationic lipids structurally related to DOTMA are described in U.S. Patent No. 4,897,355, which is herein incorporated by reference in its entirety.

TRANSFECTAM belongs to a group of cationic lipids called lipopolamines (also referred to as second-generation cationic lipids) that differ from the other lipids used in gene transfer mostly by their spermine head group. The polycationic spermine head group promotes the formation of lipoplexes with better-defined structures (*e.g.*, 50 to 100 nm) (Remy J.S. *et al.*, "Gene Transfer with Lipospermines and Polyethylenimines", *Adv. Drug Deliv. Rev.*, 1998, 30:85-95).

Another useful group of cationic lipids related to DOTMA and DOTAP that may be utilized are commonly called DORI-ethers or DORI-esters, such as (DL-1-O-oleyl-2-oleyl-3-dimethylaminopropyl- β -hydroxyethylammonium or DL-1-oleyl-2-O-oleyl-3-dimethyl-aminopropyl- β -hydroxyethylammonium). DORI lipids differ from DOTMA and DOTAP in that one of the methyl groups of the trimethylammonium group is replaced with a hydroxyethyl group. The oleoyl groups of DORI lipids can be replaced with other alkyl or alkene groups, such as palmitoyl or stearoyl groups. The hydroxyl group of the DORI-type lipids can be used as a site for further functionalization, for example for esterification to amines, like carboxyspermine. Additional cationic lipids which can be employed in the particles, compositions, and methods of the present invention include those described in International Publication No. WO 91/15501, which is herein incorporated by reference in its entirety. Cationic sterol derivatives, like 3 β [N-(N',N'-dimethylaminoethane)carbonyl] cholesterol (DC-Chol) in which cholesterol is linked to a trialkylammonium group, can also be employed in the present invention. DC-Chol is reported to provide more efficient transfection and lower toxicity than DOTMA-containing liposomes for some cell lines. DC-Chol polyamine variants such as those described in International Publication No. WO 97/45442 may also be used. Polycationic lipids containing carboxyspermine are also useful in the delivery vectors or complexes of this invention. EP-A-304111 describes carboxyspermine containing cationic lipids including 5-carboxyspermylglycine dioctadecyl-amide (DOGS), as referenced above, and dipalmitoylphosphatidylethanolamine 5-carboxyspermylamide (DPPES). Additional cationic lipids can be obtained by replacing the octadecyl and palmitoyl groups of DOGS and DPPES, respectively, with other alkyl or alkene groups. Cationic lipids can

optionally be combined with non-cationic co-lipids, preferably neutral lipids, to form the chlipids of the invention. One or more amphiphilic compounds can optionally be incorporated in order to modify the particle's surface property.

Suitable cationic lipids include esters of the Rosenthal Inhibitor (RI) (DL-2,3-distearoyloxypropyl(dimethyl)- β -hydroxyethylammoniumbromide), as described in U.S. Patent No. 5,264,618, the contents of which is hereby incorporated by reference in its entirety. These derivatives can be prepared, for example, by acyl and alkyl substitution of 3-dimethylaminopropane diol, followed by quaternization of the amino group. Analogous phospholipids can be similarly prepared.

The polynucleotides (and particles containing them) are administered and dosed in accordance with good medical practice, taking into account the clinical condition of the individual patient, the site and method of administration, scheduling of administration, patient age, sex, body weight, and other factors known to medical practitioners. The therapeutically or pharmaceutically "effective amount" for purposes herein is thus determined by such considerations as are known in the art. A therapeutically or pharmaceutically effective amount of polynucleotide (such as an IFN-encoding and/or IFN-inducible molecule-encoding polynucleotide) is that amount necessary to provide an effective amount of the polynucleotide, or the corresponding polypeptide(s) when expressed *in vivo*. An effective amount of an agent, such as a polynucleotide or non-polynucleotide agent, or particles comprising such polynucleotide or non-polynucleotide agents, can be an amount sufficient to prevent, treat, reduce and/or ameliorate the symptoms and/or underlying causes of any cell proliferation disorder, such as lung cancer. In some instances, an "effective amount" is sufficient to eliminate the symptoms of the pathologic condition and, perhaps, overcome the condition itself. In the context of the present invention, the terms "treat" and "therapy" and the like refer to alleviate, slow the progression, prophylaxis, attenuation, or cure of an existing condition. The term "prevent", as used herein, refers to putting off, delaying, slowing, inhibiting, or otherwise stopping, reducing, or ameliorating the onset of such conditions. The therapeutic methods of the invention include prevention and/or treatment of a cell proliferation disorder.

In accordance with the present invention, a suitable single dose size is a dose that is capable of preventing or alleviating (reducing or eliminating) a symptom in a patient

when administered one or more times over a suitable time period. One of skill in the art can readily determine appropriate single dose sizes for systemic administration based on the size of a mammal and the route of administration.

In one embodiment, the cells or subject to which the particles of the invention are administered is not suffering from an RNA virus infection, such as those disclosed in Mohapatra *et al.*, international publication WO 03/092618 A2 and U.S. patent publication 2004-0009152-A1, which are incorporated herein by reference in their entirety. In another embodiment, the cells or subject to which the particles of the invention are administered is not suffering from a respiratory RNA virus infection. In another embodiment, the cells or subject to which the particles of the invention are administered is not suffering from a respiratory syncytial virus (RSV) infection.

Following administration of particles to a subject, the subject's physiological condition can be monitored in various ways well known to the skilled practitioner familiar with the hallmarks of cancer progression, or alternatively by monitoring the effects of administration of the particles on the amount and/or biological activity of the interferon and/or interferon-inducible molecule *in vivo*. Optionally, the therapeutic methods of the invention include identifying a subject suffering from a cell proliferation disorder, such as lung cancer or other cancer. Identification of the subject may include medical diagnosis of the disorder by a licensed clinician.

Mammalian species which benefit from the disclosed particles, compositions, and methods include, and are not limited to, apes, chimpanzees, orangutans, humans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests, prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales.

As used herein, the term "patient", "subject", and "host" are used herein interchangeably and intended to include such human and non-human mammalian species and cells of those species. For example, the term "host" includes one or more host cells, which may be prokaryotic (such as bacterial cells) or eukaryotic cells (such as human or

non-human mammalian cells), and may be in an *in vivo* or *in vitro* state. After particles of the invention are administered to cells *in vitro*, the cells may be administered to a subject. For example, the particles of the invention can be administered to a subject's cells *ex vivo*, followed by administration of the cells to the subject. In those cases
5 wherein the polynucleotide utilized is a naturally occurring nucleic acid sequence, the polynucleotide encoding the polypeptide product can be administered to subjects of the same species or different species from which the nucleic acid sequence naturally exists, for example. When the subject is a human or the target cells are human, it is preferred that polynucleotides encoding human interferons and/or interferon-inducible molecules
10 are utilized. However, mammalian homologs may also be used, for example.

The particles of the present invention (and compositions containing them) can be administered to a subject by any route that results in delivery and/or expression of the polynucleotide (such as plasmid DNA) or delivery of other non-polynucleotide agents carried by the particles at the desired site or sites. For example, the particles can be
15 administered intravenously (I.V.), intramuscularly (I.M.), subcutaneously (S.C.), intradermally (I.D.), orally, intranasally, *etc.*

Examples of intranasal administration can be by means of a spray, drops, powder or gel and also described in U.S. Patent No. 6,489,306, which is incorporated herein by reference in its entirety. One embodiment of the present invention is the administration
20 of the invention as a nasal spray. Alternate embodiments include administration through any oral or mucosal routes such as oral, sublingual, intravaginal or intraanal administration, and even eye drops. However, other means of drug administrations such as subcutaneous, intravenous, and transdermal are well within the scope of the present invention.

25 In various embodiments, the cell proliferation disorder may be cancer of a mucous membrane, such as adenocarcinoma or other cancer of the lung, respiratory tract, stomach, epithelium, *etc.* As used herein, a "lung cancer" includes either a primary lung tumor (for example, bronchogenic carcinoma or bronchial carcinoid) or a metastasis from a primary tumor of another organ or tissue (for example, breast, colon, ovary, prostate,
30 kidney, thyroid, stomach, peritoneum, cervix, rectum, testis, bone, or melanoma).

In preferred embodiments, for cell proliferation disorders of the respiratory tract such as the lung, the particles of the invention are administered through inhalation in a

form such as an aerosol, a nebula, a mist, an atomized sample, liquid drops, *etc.* The particles are preferably delivered to the target respiratory tract tissue with a pharmacokinetic profile that results in the delivery of an effective dose of the polynucleotide carried by the particles. In preferred embodiments, at least 1%, more preferably at least 5%, even more preferably at least 10%, still more preferably at least 20%, and most preferably at least 30% or more of the administered particles preferably undergo apical to basolateral transcytosis from the pulmonary lumen.

In certain embodiments, the tumor in a subject is a primary tumor, such as that of the lung; however, the tumor in a subject may be a secondary tumor, such as a pulmonary metastasis from a primary tumor that is not of the lung. In various embodiments, the primary tumor is selected from the group consisting of a sarcoma, an adenocarcinoma, a choriocarcinoma, and a melanoma. In other embodiments, the tumor is a colon adenocarcinoma, a breast adenocarcinoma, an Ewing's sarcoma, or an osteosarcoma. For example, the primary tumor may be a renal cell carcinoma and the secondary tumor a tumor of the lung. In various embodiments, the clinical presentation of the pulmonary metastasis is a solitary metastasis, a cannonball, a lymphangitis carcinomatosa, or a pleural effusion. A "primary" tumor is the original tumor in a subject. A "secondary" tumor is a cancer that has metastasized from the organ in which it first appeared to another organ.

Cell proliferation disorders include but are not limited to solid tumors, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid and their distant metastases. Those disorders also include lymphomas, sarcomas, adenocarcinomas, and leukemias.

Cancers of any organ can be treated, such as cancers of the colon, pancreas, breast, prostate, bone, liver, kidney, lung, testes, skin, pancreas, stomach, colorectal cancer, renal cell carcinoma, hepatocellular carcinoma, melanoma, *etc.*

Examples of breast cancer include, but are not limited to, invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma *in situ*, and lobular carcinoma *in situ*. Examples of cancers of the respiratory tract include, but are not limited to, small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma and pleuropulmonary blastoma. Examples of brain cancers include, but are not limited to, brain stem and hypophthalmic glioma, cerebellar and cerebral astrocytoma,

medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumor. Tumors of the male reproductive organs include, but are not limited to, prostate and testicular cancer. Tumors of the female reproductive organs include, but are not limited to, endometrial, cervical, ovarian, vaginal, and vulvar cancer, as well as sarcoma of the uterus. Tumors of the digestive tract include, but are not limited to, anal, colon, colorectal, esophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers. Tumors of the urinary tract include, but are not limited to, bladder, penile, kidney, renal pelvis, ureter, and urethral cancers. Eye cancers include, but are not limited to, intraocular melanoma and retinoblastoma. Examples of liver cancers include, but are not limited to, hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed hepatocellular cholangiocarcinoma. Skin cancers include, but are not limited to, squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer. Head-and-neck cancers include, but are not limited to, laryngeal, hypopharyngeal, nasopharyngeal, and/or oropharyngeal cancers, and lip and oral cavity cancer. Lymphomas include, but are not limited to, AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Hodgkin's disease, and lymphoma of the central nervous system. Sarcomas include, but are not limited to, sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma. Leukemias include, but are not limited to, acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia. In addition to reducing the proliferation of tumor cells and inducing apoptosis, the particles of the invention can also cause tumor regression, *e.g.*, a decrease in the size of a tumor, or in the extent of cancer in the body.

In addition to chemotherapeutic agents, the methods and compositions of the subject invention can incorporate treatments and agents utilizing, for example, angiogenesis inhibitors (Thalidomide, Bevacizumab), Bcl-2 antisense oligonucleotides (G3139), a PSA based vaccine, a PDGF receptor inhibitor (Gleevec), microtubule stabilizers (Epothilones), and a pro-apoptotic agent (Perifosine). Thus, the particles of the invention can be administered to a subject in combination (simultaneously or consecutively) with other agents useful for treating cell proliferation disorders (including polynucleotides encoding such agents) or other disorders. Likewise, the pharmaceutical

compositions of the subject invention can include such agents (including polynucleotides encoding such agents).

The term “polynucleotide”, as used herein, refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term
5 refers only to the primary structure of the molecule. Thus, the term includes double-stranded and single-stranded DNA, as well as double-stranded and single-stranded RNA. Thus, the term includes DNA, RNA, or DNA-DNA, DNA-RNA, or RNA-RNA hybrids, or protein nucleic acids (PNAs) formed by conjugating bases to an amino acid backbone. It also includes modifications, such as by methylation and/or by capping, and unmodified
10 forms of the polynucleotide. The nucleotides may be synthetic, or naturally derived, and may contain genes, portions of genes, or other useful polynucleotides. In one embodiment, the polynucleotide comprises DNA containing all or part of the coding sequence for a polypeptide, or a complementary sequence thereof, such as interferon and/or IFN-inducible molecule. An encoded polypeptide may be intracellular, *i.e.*,
15 retained in the cytoplasm, nucleus, or in an organelle, or may be secreted by the cell. For secretion, the natural signal sequence present in a polypeptide may be retained. When the polypeptide or peptide is a fragment of a protein, a signal sequence may be provided so that, upon secretion and processing at the processing site, the desired protein will have the natural sequence. Specific examples of coding sequences of interest for use in
20 accordance with the present invention include the polypeptide-coding sequences disclosed herein. The polynucleotides may also contain, optionally, one or more expressible marker genes for expression as an indication of successful transfection and expression of the nucleic acid sequences contained therein.

According to the present invention, an isolated nucleic acid molecule or nucleic
25 acid sequence is a nucleic acid molecule or sequence that has been removed from its natural milieu. As such, “isolated” does not necessarily reflect the extent to which the nucleic acid molecule has been purified.

The terms “polypeptide” and “protein” are used interchangeably herein and indicate a molecular chain of amino acids of any length linked through peptide bonds.
30 Thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. The terms include post-translational modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. In addition, protein

fragments, analogs, mutated or variant proteins, fusion proteins and the like are included within the meaning of polypeptide.

The particles of the present invention are useful as vectors for the delivery of polynucleotides encoding interferon (such as IFN-gamma) and/or an interferon-inducible molecule (such as 2-5 AS or IRF-1) to hosts *in vitro* or *in vivo*. The term “vector” is used to refer to any molecule (*e.g.*, nucleic acid or plasmid) usable to transfer a polynucleotide, such as coding sequence information (*e.g.*, nucleic acid sequence encoding a protein or other polypeptide), to a host cell. A vector typically includes a replicon in which another polynucleotide segment is attached, such as to bring about the replication and/or expression of the attached segment. The term includes expression vectors, cloning vectors, and the like. Thus, the term includes gene expression vectors capable of delivery/transfer of exogenous nucleic acid sequences into a host cell. The term “expression vector” refers to a vector that is suitable for use in a host cell (*e.g.*, a subject’s cell, tissue culture cell, cells of a cell line, *etc.*) and contains nucleic acid sequences which direct and/or control the expression of exogenous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present. Nucleic acid sequences can be modified according to methods known in the art to provide optimal codon usage for expression in a particular expression system. The vector of the present invention may include elements to control targeting, expression and transcription of the nucleic acid sequence in a cell selective manner as is known in the art. The vector can include a control sequence, such as a promoter for controlling transcription of the exogenous material and can be either a constitutive or inducible promoter to allow selective transcription. The expression vector can also include a selection gene.

Each particle of the invention comprises a polynucleotide that is a coding sequence for an interferon, IFN-inducible molecule, or both. A “coding sequence” is a polynucleotide sequence that is transcribed into mRNA and/or translated into a polypeptide. The boundaries of the coding sequence are determined by a translation start codon at the 5’-terminus and a translation stop codon at the 3’-terminus. A coding sequence can include, but is not limited to, mRNA, cDNA, and recombinant polynucleotide sequences. Variants or analogs may be prepared by the deletion of a portion of the coding sequence, by insertion of a sequence, and/or by substitution of one

or more nucleotides within the sequence. For example, the particles of the present invention may be used to deliver coding sequences for interferon gamma, or variants or analogs thereof. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art (See, *e.g.*, Sambrook *et al.*,
5 Molecular Cloning: A Laboratory Manual, Second Edition, 1989; DNA Cloning, Vols. I and II, D.N. Glover ed., 1985). Optionally, the polynucleotides used in the particles of the present invention, and composition and methods of the invention that utilize such particles, can include non-coding sequences.

The term “operably-linked” is used herein to refer to an arrangement of flanking
10 control sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking control sequence operably-linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence under conditions compatible with the control sequences. For example, a coding sequence is operably-linked to a promoter
15 when the promoter is capable of directing transcription of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence, and the promoter sequence can still be considered “operably-linked” to the coding sequence.
20 Each nucleotide sequence coding for a polypeptide will typically have its own operably-linked promoter sequence. The promoter can be a constitutive promoter, or an inducible promoter to allow selective transcription. Optionally, the promoter can be a cell-specific or tissue-specific promoter. Promoters can be chosen based on the cell-type or tissue-type that is targeted for delivery or treatment, for example.

25 Suitable promoters include any that are known in the art or yet to be identified that will cause expression of interferon-encoding nucleic acid sequences or IFN-inducible molecule-encoding nucleic acid sequences in mammalian cells. Suitable promoters and other regulatory sequences can be selected as is desirable for a particular application. The promoters can be inducible, tissue-specific, or event-specific, as necessary. For example,
30 the cytomegalovirus (CMV) promoter (Boshart *et al.*, *Cell*, 1985, 41:521-530) and SV40 promoter (Subramani *et al.*, *Mol. Cell. Biol.*, 1981, 1:854-864) have been found to be suitable, but others can be used as well. Optionally, the polynucleotide used in the

particles of the subject invention includes a sequence encoding a signal peptide upstream of the interferon-encoding and/or IFN-inducible molecule-encoding sequence, thereby permitting secretion of the interferon and/or IFN-inducible molecule from a host cell. Also, various promoters may be used to limit the expression of the polypeptide in specific
5 cells or tissues, such as lung cells.

A tissue-specific and/or event-specific promoter or transcription element that responds to the target microenvironment and physiology can also be utilized for increased transgene expression at the desired site. There has been an immense amount of research activity directed at strategies for enhancing the transcriptional activity of weak tissue-specific promoters or otherwise increasing transgene expression with vectors. It is
10 possible for such strategies to provide enhancement of gene expression equal to one or two orders of magnitude, for example (see Nettelbeck *et al.*, *Gene Ther.*, 1998, 5(12):1656-1664 and Qin *et al.*, *Hum. Gene Ther.*, 1997, 8(17):2019-2019). Examples of cardiac-specific promoters are the ventricular form of MLC-2v promoter (see, Zhu *et al.*,
15 *Mol. Cell Biol.*, 1993, 13:4432-4444, Navankasattusas *et al.*, *Mol. Cell Biol.*, 1992, 12:1469-1479, 1992) and myosin light chain-2 promoter (Franz *et al.*, *Circ. Res.*, 1993, 73:629-638). The E-cadherin promoter directs expression specific to epithelial cells (Behrens *et al.*, *PNAS*, 1991, 88:11495-11499), while the estrogen receptor (ER) 3 gene promoter directs expression specifically to the breast epithelium (Hopp *et al.*, *J.*
20 *Mammary Gland Biol. Neoplasia*, 1998, 3:73-83). The human C-reactive protein (CRP) gene promoter (Ruther *et al.*, *Oncogene* 8:87-93, 1993) is a liver-specific promoter. An example of a muscle-specific gene promoter is human enolase (ENO3) (Peshavaria *et al.*, *Biochem. J.*, 1993, 292(Pt 3):701-704). A number of brain-specific promoters are available such as the thy-1 antigen and gamma-enolase promoters (Vibert *et al.*, *Eur. J.*
25 *Biochem.* 181:33-39, 1989). The prostate-specific antigen promoter provides prostate tissue specificity (Pang *et al.*, *Gene Ther.*, 1995, 6(11):1417-1426; Lee *et al.*, *Anticancer Res.*, 1996, 16(4A):1805-1811). The surfactant protein B promoter provides lung specificity (Strayer *et al.*, *Am. J. Respir. Cell Mol. Biol.*, 1998, 18(1):1-11). Any of the
30 aforementioned promoters may be selected for targeted or regulated expression of the interferon-encoding and/or IFN-inducible protein-encoding polynucleotide.

The particles of the present invention can be targeted through various means. As indicated above, tissue-specific promoters or event-specific promoters may be utilized

with polynucleotides encoding interferon and/or IFN-inducible molecules to further optimize and localize expression at target sites, such as within diseased tissues (*e.g.*, cancer cells or tissues containing cancer cells). Robson *et al.* review various methodologies and vectors available for delivering and expressing a polynucleotide *in vivo* for the purpose of treating cancer (Robson, T. Hirst, D.G., *J. Biomed. and Biotechnol.*, 2003, 2003(2):110-137, which is hereby incorporated by reference herein in its entirety). Among the various targeting techniques available, transcriptional targeting using tissue-specific and event-specific transcriptional control elements is discussed. For example, Table 1 at page 112 of the Robson *et al.* publication lists several tissue-specific promoters useful in cancer therapy. Tables 2-4 of the Robson *et al.* publication list tumor-specific promoters, tumor environment-specific promoters, and exogenously controlled inducible promoters, many of which were available at the time the patent application was filed. The successful delivery and expression of the p53 tumor suppressor gene *in vivo* has been documented (Horowitz, J. *Curr. Opin. Mol. Ther.*, 1999, 1(4):500-509; Von Gruenigen, V.E. *et al. Int. J. Gynecol. Cancer*, 1999, 9(5):365-372; Fujiwara, T. *et al., Mol. Urol.*, 2000, 4(2):51-54, respectively).

Many techniques for delivery of drugs and proteins are available in the art to reduce the effects of enzymatic degradation, to facilitate cell uptake, and to reduce any potential toxicity to normal (undiseased) cells, *etc.* Such methods and reagents can be utilized for administration of particles of the invention and their polynucleotide cargo to cells *in vitro* or *in vivo*. For example, peptides known as "cell penetrating peptides" (CPP) or "protein transduction domains" (PTD) have an ability to cross the cell membrane and enter the cell. PTDs can be linked to a cargo moiety such as a drug, peptide, or full-length protein, and can transport the moiety across the cell membrane. One well characterized PTD is the human immunodeficient virus (HIV)-1 Tat peptide (see, for example, Frankel *et al.*, U.S. Patent Nos. 5,804,604; 5,747,641; 6,674,980; 5,670,617; and 5,652,122; Fawell, S. *et al., Proc. Natl. Acad. Sci. U.S.A.*, 1994, 91:664-668). Peptides such as the homeodomain of *Drosophila antennapedia* (ANTp) and arginine-rich peptides display similar properties (Derossi, D. *et al., J. Biol. Chem.*, 1994, 269:10444-10450; Derossi, D. *et al., Trends Cell Biol.*, 1998, 8:84-87; Rojas, M. *et al., Nat. Biotechnol.*, 1998, 16:370-375; Futaki, S. *et al., J. Biol. Chem.*, 2001, 276:5836-5840). VP22, a tegument protein from Herpes simplex virus type 1 (HSV-1), also has the

ability to transport proteins across a cell membrane (Elliot *et al.*, *Cell*, 1997, 88:223-233; Schwarze S.R. *et al.*, *Trends Pharmacol. Sci.*, 2000, 21:45-48). A common feature of these carriers is that they are highly basic and hydrophilic (Schwarze S.R. *et al.*, *Trends Cell Biol.*, 2000, 10:290-295). Coupling of these carriers to marker proteins such as beta-galactosidase has been shown to confer efficient internalization of the marker protein into cells. More recently, chimeric, in-frame fusion proteins containing these carriers have been used to deliver proteins to a wide spectrum of cell types both *in vitro* and *in vivo*. For example, VP22-p53 chimeric protein retained its ability to spread between cells and its pro-apoptotic activity, and had a widespread cytotoxic effect in p53 negative human osteosarcoma cells *in vitro* (Phelan, A. *et al.*, *Nature Biotechnol.*, 1998, 16:440-443). Intraperitoneal injection of the beta-galactosidase protein fused to the HIV-1 Tat peptide resulted in delivery of the biologically active fusion protein to all tissues in mice, including the brain (Schwarze S.R. *et al.*, *Science*, 1999, 285:1569-1572).

Liposomes of various compositions can also be used for site-specific delivery of proteins and drugs (Witschi, C. *et al.*, *Pharm. Res.*, 1999, 16:382-390; Yeh, M.K. *et al.*, *Pharm. Res.*, 1996, 16:1693-1698). The interaction between the liposomes and their cargo usually relies on hydrophobic interactions or charge attractions, particularly in the case of cationic lipid delivery systems (Zelphati, O. *et al.*, *J. Biol. Chem.*, 2001, 276:35103-35110). Tat peptide-bearing liposomes have also been constructed and used to deliver cargo directly into the cytoplasm, bypassing the endocytotic pathway (Torchilin V.P. *et al.*, *Biochim. Biophys. Acta-Biomembranes*, 2001, 1511:397-411; Torchilin V.P. *et al.*, *Proc. Natl. Acad. Sci. USA*, 2001, 98:8786-8791). When encapsulated in sugar-grafted liposomes, pentamidine isethionate and a derivative have been found to be more potent in comparison to normal liposome-encapsulated drug or to the free drug (Banerjee, G. *et al.*, *J. Antimicrob. Chemother.*, 1996, 38(1):145-150). A thermo-sensitive liposomal taxol formulation (heat-mediated targeted drug delivery) has been administered *in vivo* to tumor-bearing mice in combination with local hyperthermia, and a significant reduction in tumor volume and an increase in survival time was observed compared to the equivalent dose of free taxol with or without hyperthermia (Sharma, D. *et al.*, *Melanoma Res.*, 1998, 8(3):240-244). Topical application of liposome preparations for delivery of insulin, IFN-alpha, IFN-gamma, and prostaglandin E1 have met with some success (Cevc G. *et al.*, *Biochim. Biophys. Acta*, 1998, 1368:201-215; Foldvari M. *et al.*, *J. Liposome*

Res., 1997, 7:115-126; Short S.M. *et al.*, *Pharm. Res.*, 1996, 13:1020-1027; Foldvari M. *et al.*, *Urology*, 1998, 52(5):838-843; U.S. Patent No. 5,853,755).

Antibodies represent another targeting device that may make particle uptake tissue-specific or cell-specific (Mastrobattista, E. *et al.*, *Biochim. Biophys. Acta*, 1999, 1419(2):353-363; Mastrobattista, E. *et al.*, *Adv. Drug Deliv. Rev.*, 1999, 40(1-2):103-127). The liposome approach offers several advantages, including the ability to slowly release encapsulated drugs and proteins, the capability of evading the immune system and proteolytic enzymes, and the ability to target tumors and cause preferentially accumulation in tumor tissues and their metastases by extravasation through their leaky neovasculature. Other carriers have also been used to deliver anti-cancer drugs to neoplastic cells, such as polyvinylpyrrolidone nanoparticles and maleylated bovine serum albumin (Sharma, D. *et al.*, *Oncol. Res.*, 1996, 8(7-8):281-286; Mukhopadhyay, A. *et al.*, *FEBS Lett.*, 1995, 376(1-2):95-98). Thus, using targeting and encapsulation technologies, which are very versatile and amenable to rational design and modification, delivery of particles of the invention to desired cells can be further facilitated.

As indicated above, the particles of the present invention can include a lipid component, such as a liposome. According to the present invention, a liposome comprises a lipid composition that is capable of fusing with the plasma membrane of a cell, thereby allowing the liposome to deliver a nucleic acid molecule and/or a protein composition into a cell. Some preferred liposomes include those liposomes commonly used in gene delivery methods known to those of skill in the art. Some preferred liposome delivery vehicles comprise multilamellar vesicle (MLV) lipids and extruded lipids, although the invention is not limited to such liposomes. Methods for preparation of MLVs are well known in the art. "Extruded lipids" are also contemplated. Extruded lipids are lipids that are prepared similarly to MLV lipids, but which are subsequently extruded through filters of decreasing size, as described in Templeton *et al.*, *Nature Biotech.*, 1997, 15:647-652, which is incorporated herein by reference in its entirety. Small unilamellar vesicle (SUV) lipids can also be used for preparing particles of the present invention. Other preferred liposome delivery vehicles comprise liposomes having a polycationic lipid composition (*i.e.*, cationic liposomes). For example, cationic liposome compositions include, but are not limited to, any cationic liposome complexed with cholesterol, and without limitation, include DOTMA and cholesterol, DOTAP and

cholesterol, DOTIM and cholesterol, and DDAB and cholesterol. Liposomes utilized in the present invention can be any size, including from about 10 to 1000 nanometers (nm), or any size in between.

A liposome delivery vehicle can be modified to target a particular site in a mammal, thereby targeting and making use of an interferon-encoding and/or IFN-inducible molecule-encoding nucleic acid molecule of the present invention at that site. Suitable modifications include manipulating the chemical formula of the lipid portion of the delivery vehicle. Manipulating the chemical formula of the lipid portion of the delivery vehicle can elicit the extracellular or intracellular targeting of the delivery vehicle. For example, a chemical can be added to the lipid formula of a liposome that alters the charge of the lipid bilayer of the liposome so that the liposome fuses with particular cells having particular charge characteristics. In some embodiments, a liposome can be directed to a particular target cell or tissue by using a targeting agent, such as an antibody, soluble receptor or ligand, incorporated with the liposome, to target a particular cell or tissue to which the targeting molecule can bind. Targeting liposomes are described, for example, in Ho *et al.*, *Biochemistry*, 1986, 25: 5500-6; Ho *et al.*, *J Biol Chem*, 1987a, 262: 13979-84; Ho *et al.*, *J Biol Chem*, 1987b, 262: 13973-8; and U.S. Patent No. 4,957,735 to Huang *et al.*, each of which is incorporated herein by reference in its entirety). In one embodiment, if avoidance of the efficient uptake of injected liposomes by reticuloendothelial system cells due to opsonization of liposomes by plasma proteins or other factors is desired, hydrophilic lipids, such as gangliosides (Allen *et al.*, *FEBS Lett*, 1987, 223: 42-6) or polyethylene glycol (PEG)-derived lipids (Klibanov *et al.*, *FEBS Lett*, 1990, 268: 235-7), can be incorporated into the bilayer of a conventional liposome to form the so-called sterically-stabilized or "stealth" liposomes (Woodle *et al.*, *Biochim Biophys Acta*, 1992, 1113: 171-99). Variations of such liposomes are described, for example, in U.S. Patent No. 5,705,187 to Unger *et al.*, U.S. Patent No. 5,820,873 to Choi *et al.*, U.S. Patent No. 5,817,856 to Tirosh *et al.*; U.S. Patent No. 5,686,101 to Tagawa *et al.*; U.S. Patent No. 5,043,164 to Huang *et al.*, and U.S. Patent No. 5,013,556 to Woodle *et al.*, all of which are incorporated herein by reference in their entireties).

The size of the particle provides another means for targeting the particles of the invention to cells or tissues. For example, relatively small particles of the invention can be made to efficiently target ischemic tissue and tumor tissue, as described in U.S. Patent

No. 5,527,538, and U.S. Patent Nos. 5,019,369, 5,435,989 and 5,441,745, the contents of which are hereby incorporated by reference in their entirety.

The particles of the invention can be targeted according to the mode of administration. For example, lung or other respiratory epithelial tissue can be targeted by intranasal administration, cervical cells can be targeted by intravaginal administration, and prostate tumors can be targeted by intrarectal administration. Skin cancer can be targeted by topical administration. Depending on location, tumors can be targeted locally, such as by injection, into the tumor mass.

Particles of the invention can be targeted by incorporating a ligand such as an antibody, a receptor, or other compound known to target particles such as liposomes or other vesicles to various sites. The ligands can be attached to cationic lipids used to form the particles of the present invention, or to a neutral lipid such as cholesterol used to stabilize the particle. Ligands that are specific for one or more specific cellular receptor sites are attached to a particle to form a delivery vehicle that can be targeted with a high degree of specificity to a target cell population of interest.

Suitable ligands for use in the present invention include, but are not limited to, sugars, proteins such as antibodies, hormones, lectins, major histocompatibility complex (MHC), and oligonucleotides that bind to or interact with a specific site. An important criteria for selecting an appropriate ligand is that the ligand is specific and is suitably bound to the surface of the particles in a manner which preserves the specificity. For example, the ligand can be covalently linked to the lipids used to prepare the particles. Alternatively, the ligand can be covalently bound to cholesterol or another neutral lipid, where the ligand-modified cholesterol is used to stabilize the lipid monolayer or bilayer.

The terms "transfection" and "transformation" are used interchangeably herein to refer to the insertion of an exogenous polynucleotide into a host, irrespective of the method used for the insertion, the molecular form of the polynucleotide that is inserted, or the nature of the host (*e.g.*, prokaryotic or eukaryotic). The insertion of a polynucleotide per se and the insertion of a plasmid or vector comprising the exogenous polynucleotide are included. The exogenous polynucleotide may be directly transcribed and translated by the host or host cell, maintained as a nonintegrated vector, for example, a plasmid, or alternatively, may be stably integrated into the host genome. The terms "administration"

and “treatment” are used herein interchangeably to refer to transfection of hosts *in vitro* or *in vivo*, using particles of the present invention.

The term “wild-type” (WT), as used herein, refers to the typical, most common or conventional form as it occurs in nature.

5 Thus, the present invention includes methods of gene therapy whereby polynucleotides encoding the desired gene product (an interferon, such as interferon-gamma, an IFN-inducible molecule, or both) are delivered to a subject, and the polynucleotide is expressed *in vivo*. The term “gene therapy”, as used herein, includes the transfer of genetic material (*e.g.*, polynucleotides) of interest into a host to treat or
10 prevent a genetic or acquired disease or condition phenotype, or to otherwise express the genetic material such that the encoded product is produced within the host. The genetic material of interest can encode a product (*e.g.*, a protein, polypeptide, peptide, or functional RNA) whose production *in vivo* is desired. For example, in addition to interferon and/or an IFN-inducible molecule, the genetic material can encode a hormone,
15 receptor, enzyme, polypeptide or peptide, of therapeutic and/or diagnostic value. For a review see, in general, the text “Gene Therapy” (*Advances in Pharmacology* 40, Academic Press, 1997). The genetic material may encode a product normally found within the species of the intended host, or within a different species. For example, if the polynucleotide encodes interferon-gamma, the cytokine may be human interferon-
20 gamma, or that of another mammal, for example, regardless of the intended host. Preferably, the polynucleotide encodes a product that is normally found in the species of the intended host. Alternatively, the genetic material may encode a novel product.

Two basic approaches to gene therapy have evolved: (1) *ex vivo* and (2) *in vivo* gene therapy. The methods of the subject invention encompass either or both. In *ex vivo*
25 gene therapy, host cells are removed from a patient and, while being cultured, are treated *in vitro*. Generally, a functional replacement gene is introduced into the cell via an appropriate gene delivery vehicle/method (transfection, transduction, homologous recombination, *etc.*) and an expression system as needed and then the modified cells are expanded in culture and returned to the host/patient.

30 In *in vivo* gene therapy, target host cells are not removed from the subject, rather the gene to be transferred is introduced into the cells of the recipient organism *in situ*, that

is within the recipient. Alternatively, if the host gene is defective, the gene is repaired *in situ*.

The particle of the present invention is capable of delivery/transfer of heterologous nucleic acid sequences into a prokaryotic or eukaryotic host cell *in vitro* or
5 *in vivo*. The particle may include elements to control targeting, expression and transcription of the nucleic acid sequence in a cell selective manner as is known in the art. It should be noted that often the 5'UTR and/or 3'UTR of the gene may be replaced by the 5'UTR and/or 3'UTR of other expression vehicles.

Optionally, the particles of the invention may have biologically active agents other
10 than polynucleotides as a component of the complex (either instead of, or in addition to, polynucleotides). Such biologically active agents include, but are not limited to, substances such as proteins, polypeptides, antibodies, antibody fragments, lipids, carbohydrates, and chemical compounds such as pharmaceuticals. The substances can be therapeutic agents, diagnostic materials, and/or research reagents.

The present invention includes pharmaceutical compositions comprising an
15 effective amount of particles of the invention and a pharmaceutically acceptable carrier. The pharmaceutical compositions of the subject invention can be formulated according to known methods for preparing pharmaceutically useful compositions. As used herein, the phrase "pharmaceutically acceptable carrier" means any of the standard pharmaceutically
20 acceptable carriers. The pharmaceutically acceptable carrier can include diluents, adjuvants, and vehicles, as well as implant carriers, and inert, non-toxic solid or liquid fillers, diluents, or encapsulating material that does not react with the active ingredients of the invention. Examples include, but are not limited to, phosphate buffered saline, physiological saline, water, and emulsions, such as oil/water emulsions. The carrier can
25 be a solvent or dispersing medium containing, for example, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils.

The pharmaceutically acceptable carrier can be one adapted for a particular route of administration. For example, if the particles of the present invention are intended to be
30 administered to the respiratory epithelium, a carrier appropriate for oral or intranasal administration can be used.

Formulations containing carriers are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E.W., 1995, Easton Pennsylvania, Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the subject invention. Formulations suitable for parenteral administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, *etc.* It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

As used herein, the terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include breast cancer, prostate cancer, colon cancer, squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, *e.g.*, hepatic carcinoma, bladder cancer, colorectal cancer, endometrial carcinoma, kidney cancer, and thyroid cancer.

Other non-limiting examples of cancers are basal cell carcinoma, biliary tract cancer; bone cancer; brain and CNS cancer; choriocarcinoma; connective tissue cancer; esophageal cancer; eye cancer; cancer of the head and neck; gastric cancer; intra-epithelial neoplasm; larynx cancer; lymphoma including Hodgkin's and Non-Hodgkin's lymphoma; melanoma; myeloma; neuroblastoma; oral cavity cancer (*e.g.*, lip, tongue, mouth, and pharynx); peritoneal cancer; retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; sarcoma; skin cancer; stomach cancer; testicular

cancer; uterine cancer; cancer of the urinary system, as well as other carcinomas and sarcomas.

As used herein, the term “tumor” refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. For example, a particular cancer may be characterized by a solid mass tumor. The solid tumor mass, if present, may be a primary tumor mass. A primary tumor mass refers to a growth of cancer cells in a tissue resulting from the transformation of a normal cell of that tissue. In most cases, the primary tumor mass is identified by the presence of a cyst, which can be found through visual or palpation methods, or by irregularity in shape, texture or weight of the tissue. However, some primary tumors are not palpable and can be detected only through medical imaging techniques such as X-rays (*e.g.*, mammography), or by needle aspirations. The use of these latter techniques is more common in early detection. Molecular and phenotypic analysis of cancer cells within a tissue will usually confirm if the cancer is endogenous to the tissue or if the lesion is due to metastasis from another site.

As used herein, the term “metastasis” refers to the process by which cancer cells are spread to distant parts of the body, such as from one organ and/or tissue to another not directly connected with it. The term is also used herein to refer to a tumor that develops through the metastatic process. Thus, as used herein, the term “metastasis” refers to neoplastic cell growth (*e.g.*, tumor cell growth) in an unregulated fashion and spread to distal tissues and organs of the body. As used herein, the phrase “inhibiting metastasis” refers to the particles slowing and/or preventing metastasis or the spread of neoplastic cells to a site remote from the primary growth area.

The term “anti-cancer activity”, in reference to the particles of the invention, is intended to mean an activity which is able to substantially inhibit, slow, interfere, suppress, prevent, delay and/or arrest a cancer and/or a metastasis thereof (such as initiation, growth, spread, and/or progression thereof of such cancer and/or metastasis).

As used herein, the term “growth inhibitory amount” refers to an amount which inhibits growth of a target cell, such as a tumor cell, either *in vitro* or *in vivo*, irrespective of the mechanism by which cell growth is inhibited. In a preferred embodiment, the growth inhibitory amount inhibits growth of the target cell in cell culture by greater than

about 20%, preferably greater than about 50%, most preferably greater than about 75% (*e.g.*, from about 75% to about 100%).

The therapeutic methods of the invention can be advantageously combined with at least one additional therapeutic technique, including but not limited to chemotherapy, radiation therapy, surgery (*e.g.*, surgical excision of cancerous or pre-cancerous cells), or any other therapy known to those of skill in the art of the treatment and management of cancer, such as administration of an anti-cancer agent.

As used herein, the term “anti-cancer agent” refers to a substance or treatment that inhibits the function of cancer cells, inhibits their formation, and/or causes their destruction *in vitro* or *in vivo*. Examples include, but are not limited to, cytotoxic agents (*e.g.*, 5-fluorouracil, TAXOL) and anti-signaling agents (*e.g.*, the PI3K inhibitor LY).

As used herein, the term “cytotoxic agent” refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells *in vitro* and/or *in vivo*. The term is intended to include radioactive isotopes (*e.g.*, At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², and radioactive isotopes of Lu), chemotherapeutic agents, toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, and antibodies, including fragments and/or variants thereof.

As used herein, the term “chemotherapeutic agent” is a chemical compound useful in the treatment of cancer, such as, for example, taxanes, *e.g.*, paclitaxel (TAXOL, BRISTOL-MYERS SQUIBB Oncology, Princeton, N.J.) and doxorubicin (DOXIL, Rhone-Poulenc Rorer, Antony, France), chlorambucil, vincristine, vinblastine, anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston), and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin, *etc.*

As used herein, the term “anti-signaling agent” refers to agents that interfere with cancer cell malignancy by inhibiting specific aberrant signal transduction circuits in the cell *in vitro* and/or *in vivo*. The PI3K inhibitor LY is an example of an anti-signalling agent.

The terms “comprising”, “consisting of” and “consisting essentially of” are defined according to their standard meaning. The terms may be substituted for one

another throughout the instant application in order to attach the specific meaning associated with each term.

As used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise.

5 Thus, for example, a reference to “a particle” includes more than one such particle, a reference to “a polynucleotide” includes more than one such polynucleotide, a reference to “a polypeptide” includes more than one such polypeptide, a reference to “a host cell” includes more than one such host cell, a reference to an interferon or IFN-inducible molecule includes more than one such interferon or IFN-inducible molecule, and so forth.

10 Standard molecular biology techniques known in the art and not specifically described were generally followed as in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York (1989), and in Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Md. (1989) and in Perbal, A Practical Guide to Molecular Cloning, John Wiley & Sons,
15 New York (1988), and in Watson *et al.*, Recombinant DNA, Scientific American Books, New York and in Birren *et al.* (eds) Genome Analysis: A Laboratory Manual Series, Vols. 1-4 Cold Spring Harbor Laboratory Press, New York (1998) and methodology as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659; and 5,272,057; and incorporated herein by reference. Polymerase chain reaction (PCR) was carried out
20 generally as in PCR Protocols: A Guide To Methods And Applications, Academic Press, San Diego, Calif. (1990). *In situ* (In-cell) PCR in combination with Flow Cytometry can be used for detection of cells containing specific DNA and mRNA sequences (Testoni *et al.*, *Blood*, 1996, 87:3822).

25 It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

30 Example 1—pIFN- γ induces apoptosis of HEp-2 carcinoma cells

To determine the effect of overexpression of pIFN- γ on proliferation of A549 lung epithelial cells, cells were transfected with either pIFN- γ or empty vector, pVAX

(control). Cell cycle analysis was performed using propidium iodide (PI) staining and flow cytometry 48 hours after transfection. No significant difference was observed between control and pIFN- γ -transfected cells in S1, Go-G1 and G2-M stages of the cell cycle (data not shown). However, an analysis of apoptosis using fluorescence microscopy cells transfected with pIFN- γ exhibited significantly higher apoptosis compared to cells transfected with either the control plasmid or a plasmid encoding pVAX (shown in Figure 1).

Cells were seeded into 4-well slide dishes at 104 cells per well and allowed to grow to 75% confluency. Cells were treated for 20 hours with 1000 U/ml IFN- γ . After 24 hours, cells were fixed with 4% paraformaldehyde in PBS for 25 minutes at 4°C and then permeabilized. Apoptotic cells were identified using a fluorescein-based, terminal nucleotidyl end-labeling kit (PROMEGA TUNEL Apoptosis Assay) that adds fluorescein conjugated dUTP to the 3'-hydroxyl ends of DNA fragments arising from apoptosis. After the reaction, the cells were rinsed in 2x saline citrate buffer and the nuclei were stained with DAPI. Stained cells were examined by immuno-fluorescence microscopy to determine the extent of apoptosis.

Figure 2 demonstrates the detection of p27kip expression and PARP cleavage in IFN-gamma treated HEp-2 cells. Total cell extracts of HEp-2 cells (1×10^6) treated as above were prepared after 24 and 48 hours of treatment and proteins were subjected to SDS-PAGE and western blotting was done with a monoclonal antibody to p27kip1 (A) or an antibody to PARP (B). The lanes are as follows: 1) Untreated cells, 2) IFN-gamma 100u/ml, 3) IFN-gamma 1000u/ml, 4) IFN-beta 100u/ml, 5) IFN-beta 1000u/ml, 6) untreated cells, 7) IFN-gamma 100u/ml, 8) IFN-gamma 1000u/ml, 9) IFN-beta 100u/ml, and 10) IFN-beta 1000u/ml.

The apoptosis was confirmed by analysis of PARP cleavage in these cells 48 hours after transfection, which was significantly more prominent in pIFN- γ transfected cells (Figure 2). Thus, pIFN- γ induces apoptosis of lung adenocarcinoma cells. Together, these studies indicate that pIFN- γ is an inducer of apoptosis in A549 lung adenocarcinoma cells.

Example 2—Microarray analysis of Chitosan –pIFNgamma treated lungs

Using MU11KsubA and B chips (Affymetrix), which contain probes interrogating about 11,000 murine genes and ESTs (Unigene, Build-4), as well as EST clusters from TIGR (1.0 Beta), we have identified a total of 126 differentially expressed genes whose expression level is altered in the CIN treated mouse lung in the range of –10.6- to 152.4-fold. A noteworthy observation is the up-regulation of the expression of a number of IFN-inducible genes, immune response related genes, and genes involved in signal transduction, including STAT1 and STAT4.

10 **Table 1. GENE EXPRESSION ANALYSIS IN BALB/c LUNG BY MICROARRAY**

Category of Genes	Max. Fold Change	Genes
IFN-regulated	12	IFN-induced 15 KDa protein, IFN-activable protein 204, Eukaryotic initiation factor 5, Mx protein, MIG, IP-10, Interferon regulatory factor (mirf7), interferon-activatable protein, IFN-induced protein 6-16 precursor, IFN-induced guanylate-binding protein, HLA- associated protein i (phapi) 2'-5' oilgo A synthetase
Immune-related	8.4	T-cell specific protein, RegIII gamma protein, MHC class II, MIP, Down regulatory protein (rpt-1r) of interleukin-2, T-cell receptor alpha-chain precursor, Immune-responsive gene 1 (Irg1), High affinity IgG receptor, MHC, class III antigene factor B, MEL-14, Lymphotoxin-beta, C-11, Rantes, MAMA Serine protease, Proteasome subunit (Imp7)
Signal transduction	10.8*	PDGF, GTPase IGTP, Glucocorticoid-attenuated response gene 16, Stat1, purine nucleotide binding protein, G-protein-like LRG-47, ras-related protein ora2, GTP binding protein (IRG-47), Stat 4, cathepsin s precursor, Oct binding factor 1 (OBF-1), FYN binding protein, High mobility group 2

The RNA was isolated from BALB/c lungs following 5 days of CIN treatment. The mouse chips A and B, a total of 11,000 genes, were scanned. The asterisk indicates the fold increase was uncertain, as no expression was observed in control lungs. The genes are listed in no particular order.

Example 3—Chitosan-conjugated pIFN- γ plasmid prevents metastasis of lung tumors in nude mice

BALB/c nude mice were injected intravenously with 5×10^6 A549 cells, then treated one day afterwards and at weekly intervals with pIFN- γ or control plasmid. After 4 weeks, mice were examined for lung histology. The control animals showed tumors, whereas no tumors were seen in the pIFN- γ -treated group (Figure 3). These results indicate that pIFN- γ has the potential to decrease tumor metastasis.

The results indicated in Figure 3 were obtained when BALB/c nude mice were injected with A549 cells (5×10^6 cells/mouse) intravenously and one group treated with pIFN-gamma and another group with pVAX as control. The lungs of control mice showed numerous lung nodules in contrast to mice treated with pIFN-gamma, which showed very few tumors. The lungs were removed from mice treated with nanoparticles carrying empty plasmid pVAX (control) or with pIFN-gamma (Rx) and H & E stained. The lungs of control mice showed numerous lung nodules with typical tumor cell morphology in contrast to mice treated with pIFN-gamma, which showed very few tumors.

Example 4—Development of thermogel from modified chitosan that provides sustained release

Using depolymerization methods, four novel chitosan derivative was synthesized. The products were separated by capillary gel electrophoresis. The plot shows the separation of 2 low molecular weight components (Figure 4A). Nanogene-042(NG042) is a unique low molecular weight chitosan-based carrier, which has a particle size of 155 nm (major peak, with some aggregates at 335 nm), a zeta potential of about +20 mV with typical oligomeric structure, as identified by atomic force microscopy, and significant heat-stable properties for gene transfer, with both *in vitro* and *in vivo* expression (Figures 4B and 4C). Lyophilized and resuspended NG042 particles retain functionality at ambient temperatures of 23° to 55° C. Nanogene complexes of pGL3 (firefly luciferase, Promega) was lyophilized, reconstituted with water and treated for 24 hours at RT (23° C), 42° C, 55° C, and -20° C. A549 cells were plated and transfected with the above complexes. Uptake and expression of DNA was allowed to occur for 24 hours. Luciferase activity

was determined by using Promega's Dual Assay kit. Readings were normalized to relative luminiscence units (RLU) per mg protein.

Another carrier, Nanogene-044 (NG044), is soluble in water and supports sustained gene expression *in vivo* (Figure 5A). It also exhibits thermo-gelling properties, *i.e.*, it is liquid at room temperature and forms a gel at temperatures above 37°C (Figure 5B). NG045 is a 1000-dalton oligomeric chitosan that is water soluble and shows sustained drug delivery following

NG044 was found to form a gel upon reacting with 2-glycerol phosphate, while NG042, another depolymerized chitosan does not.

To establish length of gene expression, Nanogene 044 (NG044) particles were complexed with DNA (5:1) encoding green fluorescent protein and a hydrogel was formed. The hydrogel was administered intranasally to groups of mice (n=4). Mice were sacrificed on the indicated days and broncho-alveolar lavage cells were examined by fluorescent microscopy. Another group received NG044 with pEGFP without gelling (Control). Gene expression in the mouse lung was measured by EGFP expression in BAL cells 10 and 20 days after administration. The results at day 10 were similar (not shown) for control and hydrogel, whereas after 20 days, mice given hydrogel continued EGFP show expression and no expression was detected in control mice.

Overall, the notion of intranasal chitosan nanoparticles carrying pIFN- γ for the treatment of cancer is based on the preliminary results that pIFN- γ may induce epithelial cell production of NO, which is known to possess anti-tumor effects, apoptosis of carcinoma cells, and abrogation of lung nodule formation in a murine model of lung metastasis. It will be seen that the objects set forth above, and those made apparent from the foregoing description, are efficiently attained and since certain changes may be made in the above construction without departing from the scope of the invention, it is intended that all matters contained in the foregoing description or shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

All patents, patent applications, provisional applications, and publications (including information associated with sequence accession numbers) referred to or cited herein are incorporated by reference in their entirety, including all figures and tables, to the extent they are not inconsistent with the explicit teachings of this specification.

CLAIMS

I claim:

1. A method for treating a cell proliferation disorder in a subject, comprising administering a therapeutically effective amount of particles to the subject, wherein the particles comprise: a polynucleotide encoding an interferon, an interferon-inducible molecule, or both; and a chitin-containing component associated with the polynucleotide, wherein the polynucleotide is expressed in the subject and cell proliferation is reduced.
2. The method of claim 1, wherein the interferon is selected from the group consisting of alpha interferon, beta interferon, gamma interferon, omega interferon, and lambda interferon, or a biologically active fragment or derivative thereof.
3. The method of claim 1, wherein the interferon is gamma interferon.
4. The method of claim 1, wherein the interferon is a hybrid interferon.
5. The method of claim 1, wherein the interferon inducible molecule comprises interferon regulatory factor-1 (IRF-1).
6. The method of claim 1, wherein the interferon-inducible molecule comprises 2'-5' oligoadenylate synthetase, interferon regulatory factor-1 (IRF-1), or both.
7. The method of claim 1, wherein the interferon-inducible molecule comprises a catalytically active subunit of 2'-5' oligoadenylate synthetase selected from the group consisting of p40, p69, and p100 subunit.
8. The method of claim 1, wherein the 2'-5' oligoadenylate synthetase comprises at least one splice variant selected from the group consisting of 40kDa, 42kDa, 46kDa, 69 kDa, and 71 kDa.

9. The method of claim 1, wherein the chitin-containing component comprises chitosan or a chitosan derivative.

10. The method of claim 1, wherein the particles further comprise a lipid component associated with the chitin-containing component and the polynucleotide.

11. The method of claim 1, wherein the cell proliferation disorder is a cancer of the respiratory tract.

12. The method of claim 1, wherein the cell proliferation disorder is lung cancer.

13. The method of claim 1, wherein the particles are administered to the subject via a mucosal route.

14. The method of claim 1, wherein the particles are administered to the subject intranasally.

15. The method of claim 1, wherein the particles are administered to the subject as a spray, drops, powder, gel, or a combination of two or more of the foregoing.

16. The method of claim 1, wherein the subject is human.

17. The method of claim 1, wherein the subject is suffering from a cell proliferation disorder.

18. The method of claim 1, wherein the subject has been diagnosed with the cell proliferation disorder prior to said administering.

19. A method of inducing apoptosis in a cancer cell, comprising contacting a target cancer cell *in vitro* or *in vivo* with an effective amount of particles comprising: a polynucleotide encoding an interferon, an interferon-inducible molecule, or both; and a

chitin-containing component associated with the polynucleotide, wherein the polynucleotide is expressed in the cancer cell and apoptosis is induced.

20. The method of claim 19, wherein the interferon is selected from the group consisting of alpha interferon, beta interferon, gamma interferon, omega interferon, and lambda interferon, or a biologically active fragment or derivative thereof.

21. The method of claim 19, wherein the interferon-inducible molecule comprises 2'-5' oligoadenylate synthetase, interferon regulatory factor-1 (IRF-1), or both.

22. The method of claim 19, wherein the cancer cell is a respiratory epithelial cell.

23. A particle comprising a polynucleotide encoding an interferon, an interferon-inducible molecule, or both; and a chitin-containing component associated with the polynucleotide.

24. The particle of claim 23, wherein said chitin-containing component comprises chitosan or a chitosan derivative.

25. The particle of claim 23, wherein said particle further comprises a lipid component associated with the chitin-containing component and the polynucleotide.

26. The particle of claim 23, wherein said particle comprises a polynucleotide encoding said interferon and said interferon-inducible molecule.

27. The particle of claim 23, wherein said particle comprises a polynucleotide encoding said interferon and 2'-5' oligoadenylate synthetase.

28. The particle of claim 23, wherein said particle comprises a polynucleotide encoding said interferon and interferon regulatory factor-1 (IRF-1).

29. The particle of claim 23, wherein said particle comprises a polynucleotide encoding 2'-5' oligoadenylate synthetase and interferon regulatory factor-1 (IRF-1).

30. The particle of claim 23, wherein said particle comprises a polynucleotide encoding said interferon, 2'-5' oligoadenylate synthetase, and interferon regulatory factor-1 (IRF-1).

31. The particle of claim 23, wherein said polynucleotide comprises one or more nucleotide sequences selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, and 31.




FIG. 1

2/9




FIG. 2




FIG. 3

3/9




FIG. 4A

Size & Zeta analysis




FIG. 4B

AFM analysis



FIG. 4C

6/9




FIG. 4D




FIG. 5A

Thermogelling

NG042

NG044




FIG. 5B




FIG. 5C

SEQUENCE LISTING

<110> University of South Florida
Mohapatra, Shyam

<120> Nanogene Therapy For Cell Proliferation Disorders

<130> USF-216PTCZ

<150> 60/565,756
<151> 2004-04-27

<160> 31

<170> PatentIn version 3.3

<210> 1
<211> 1203
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(1203)

<300>
<301> Aissouni, Y., et al.
<302> The cleavage/polyadenylation activity triggered by a U-rich motif
sequence
<303> J. Biol. Chem.
<304> 277
<305> 39
<306> 35808-35814
<307> 2002
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Behera, A.K., et al.
<302> 2'-5' Oligoadenylate synthesis plays a critical role in
interferon-gamma inhibition
<303> J. Biol. Chem.
<304> 277
<305> 28
<306> 25601-25608
<307> 2002
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Sarker, S.N., et al.
<302> Identification of the substrate-binding sites of
2'-5'-oligoadenylate synthetase
<303> J. Biol. Chem.
<304> 277
<305> 27
<306> 24321-24330
<307> 2002

<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Hovnanian, A., et al.
<302> The human 2',5'-oligoadenylate synthetase locus is composed of
three distinct genes
<303> Genomics
<304> 52
<305> 3
<306> 267-277
<307> 1998
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Renault, B., et al.
<302> A sequence-ready physical map of a region of 12q24.1
<303> Genomics
<304> 45
<305> 2
<306> 271-278
<307> 1997
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Nechiporuk, T., et al.
<302> A high-resolution PAC and BAC map of the SCA2 region
<303> Genomics
<304> 44
<305> 3
<306> 321-329
<307> 1997
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Wathelet, M.G., et al.
<302> Cloning and chromosomal location of human genes inducible by type
I interferon
<303> Somat. Cell Mol. Genet.
<304> 14
<305> 5
<306> 415-426
<307> 1988
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Rutherford, M.N., et al.
<302> Interferon-induced binding of nuclear factors to promoter
elements of the 2-5A synthetase gene
<303> EMBO J.
<304> 7
<305> 3
<306> 751-759
<307> 1988
<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Wathelet, M.G., et al.

<302> New inducers revealed by the promoter sequence analysis of two interferon-activated human genes

<303> Eur. J. Biochem.

<304> 169

<305> 2

<306> 313-321

<307> 1987

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Benech, P., et al.

<302> Interferon-responsive regulatory elements in the promoter of the human 2',5'-oligo(A) synthetase gene

<303> Mol. Cell. Biol.

<304> 7

<305> 12

<306> 4498-4504

<307> 1987

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Hovanessian, A.G., et al.

<302> Identification of 69-kd and 100-kd forms of 2-5A synthetase

<303> EMBO J.

<304> 6

<305> 5

<306> 1273-1280

<307> 1987

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Williams, B.R., et al.

<302> Interferon-regulated human 2-5A synthetase gene maps to chromosome

<303> Somat. Cell Mol. Genet.

<304> 12

<305> 4

<306> 403-408

<307> 1986

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Shiojiri, S., et al.

<302> Structure and expression of a cloned cDNA for human (2'-5') oligoadenylate synthetase

<303> J. Biochem.

<304> 99

<305> 5

<306> 1455-1464

<307> 1986

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Wathelet, M., et al.

<302> Full-length sequence and expression of the 42 kDa 2-5A synthetase

<303> FEBS Lett.

<304> 196

<305> 1

<306> 113-120

<307> 1986

<308> NCBI/NM_016816

<309> 2003-04-06

<300>

<301> Benech, P., et al.

<302> Structure of two forms of the interferon-induced (2'-5') oligo A synthetase of human cells

<303> EMBO J.

<304> 4

<305> 9

<306> 2249-2256

<307> 1985

<308> NCBI/NM 016816

<309> 2003-04-06

<300>

<301> Saunders, M.E., et al.

<302> Human 2-5A synthetase: characterization of a novel cDNA and corresponding gene structure

<303> EMBO J.

<304> 4

<305> 7

<306> 1761-1768

<307> 1985

<308> NCBI/NM 016816

<309> 2003-04-06

<300>

<301> Merlin, G., et al.

<302> Molecular cloning and sequence of partial cDNA for
interferon-induced (2'-5')oligo(A_ synthetase mRNA from human
cells

<303> Proc. Natl. Acad. Sci. U.S.A.

<304> 80

<305> 16

<306> 4904-4908

<307> 1983

<308> NCBI/NM 016816

<309> 2003-04-06

<400> 1

atg	atg	gat	ctc	aga	aat	acc	cca	gcc	aaa	tct	ctg	gac	aag	ttc	att	48
Met	Met	Asp	Leu	Arg	Asn	Thr	Pro	Ala	Lys	Ser	Leu	Asp	Lys	Phe	Ile	
1				5					10					15		

gaa gac tat ctc ttg cca gac acg tgt ttc cgc atg caa atc gac cat 96
Glu Asp Tyr Leu Leu Pro Asp Thr Cys Phe Arg Met Gln Ile Asp His
20 25 30

gcc att gac atc atc tgt ggg ttc ctg aag gaa agg tgc ttc cga ggt Ala Ile Asp Ile Ile Cys Gly Phe Leu Lys Glu Arg Cys Phe Arg Gly 35 40 45	144
agc tcc tac cct gtg tgt gtg tcc aag gtg gta aag ggt ggc tcc tca Ser Ser Tyr Pro Val Cys Val Ser Lys Val Val Lys Gly Gly Ser Ser 50 55 60	192
ggc aag ggc acc acc ctc aga ggc cga tct gac gct gac ctg gtt gtc Gly Lys Gly Thr Thr Leu Arg Gly Arg Ser Asp Ala Asp Leu Val Val 65 70 75 80	240
ttc ctc agt cct ctc acc act ttt cag gat cag tta aat cgc cgg gga Phe Leu Ser Pro Leu Thr Thr Phe Gln Asp Gln Leu Asn Arg Arg Gly 85 90 95	288
gag ttc atc cag gaa att agg aga cag ctg gaa gcc tgt caa aga gag Glu Phe Ile Gln Glu Ile Arg Arg Gln Leu Glu Ala Cys Gln Arg Glu 100 105 110	336
aga gca ctt tcc gtg aag ttt gag gtc cag gct cca cgc tgg ggc aac Arg Ala Leu Ser Val Lys Phe Glu Val Gln Ala Pro Arg Trp Gly Asn 115 120 125	384
ccc cgt gcg ctc agc ttc gta ctg agt tcg ctc cag ctc ggg gag ggg Pro Arg Ala Leu Ser Phe Val Leu Ser Ser Leu Gln Leu Gly Glu Gly 130 135 140	432
gtg gag ttc gat gtg ctg cct gcc ttt gat gcc ctg ggt cag ttg act Val Glu Phe Asp Val Leu Pro Ala Phe Asp Ala Leu Gly Gln Leu Thr 145 150 155 160	480
ggc agc tat aaa cct aac ccc caa atc tat gtc aag ctc atc gag gag Gly Ser Tyr Lys Pro Asn Pro Gln Ile Tyr Val Lys Leu Ile Glu Glu 165 170 175	528
tgc acc gac ctg cag aaa gag ggc gag ttc tcc acc tgc ttc aca gaa Cys Thr Asp Leu Gln Lys Glu Gly Glu Phe Ser Thr Cys Phe Thr Glu 180 185 190	576
cta cag aga gac ttc ctg aag cag cgc ccc acc aag ctc aag agc ctc Leu Gln Arg Asp Phe Leu Lys Gln Arg Pro Thr Lys Leu Lys Ser Leu 195 200 205	624
atc cgc cta gtc aag cac tgg tac caa aat tgt aag aag aag ctt ggg Ile Arg Leu Val Lys His Trp Tyr Gln Asn Cys Lys Lys Lys Leu Gly 210 215 220	672
aag ctg cca cct cag tat gcc ctg gag ctc ctg acg gtc tat gct tgg Lys Leu Pro Pro Gln Tyr Ala Leu Glu Leu Leu Thr Val Tyr Ala Trp 225 230 235 240	720
gag cga ggg agc atg aaa aca cat ttc aac aca gcc caa gga ttt cgg Glu Arg Gly Ser Met Lys Thr His Phe Asn Thr Ala Gln Gly Phe Arg 245 250 255	768
acg gtc ttg gaa tta gtc ata aac tac cag caa ctc tgc atc tac tgg Thr Val Leu Glu Leu Val Ile Asn Tyr Gln Gln Leu Cys Ile Tyr Trp 260 265 270	816

aca aag tat tat gac ttt aaa aac ccc att att gaa aag tac ctg aga 864
 Thr Lys Tyr Tyr Asp Phe Lys Asn Pro Ile Ile Glu Lys Tyr Leu Arg
 275 280 285
 agg cag ctc acg aaa ccc agg cct gtg atc ctg gac ccg gcg gac cct 912
 Arg Gln Leu Thr Lys Pro Arg Pro Val Ile Leu Asp Pro Ala Asp Pro
 290 295 300
 aca gga aac ttg ggt ggt gga gac cca aag ggt tgg agg cag ctg gca 960
 Thr Gly Asn Leu Gly Gly Gly Asp Pro Lys Gly Trp Arg Gln Leu Ala
 305 310 315 320
 caa gag gct gag gcc tgg ctg aat tac cca tgc ttt aag aat tgg gat 1008
 Gln Glu Ala Glu Ala Trp Leu Asn Tyr Pro Cys Phe Lys Asn Trp Asp
 325 330 335
 ggg tcc cca gtg agc tcc tgg att ctg ctg gct gaa agc aac agt aca 1056
 Gly Ser Pro Val Ser Ser Trp Ile Leu Leu Ala Glu Ser Asn Ser Thr
 340 345 350
 gac gat gag acc gac gat ccc agg acg tat cag aaa tat ggt tac att 1104
 Asp Asp Glu Thr Asp Asp Pro Arg Thr Tyr Gln Lys Tyr Gly Tyr Ile
 355 360 365
 gga aca cat gag tac cct cat ttc tct cat aga ccc agc acg ctc cag 1152
 Gly Thr His Glu Tyr Pro His Phe Ser His Arg Pro Ser Thr Leu Gln
 370 375 380
 gca gca tcc acc cca cag gca gaa gag gac tgg acc tgc acc atc ctc 1200
 Ala Ala Ser Thr Pro Gln Ala Glu Glu Asp Trp Thr Cys Thr Ile Leu
 385 390 395 400
 tga 1203

<210> 2
 <211> 400
 <212> PRT
 <213> Homo sapiens

<400> 2

Met Met Asp Leu Arg Asn Thr Pro Ala Lys Ser Leu Asp Lys Phe Ile
 1 5 10 15

Glu Asp Tyr Leu Leu Pro Asp Thr Cys Phe Arg Met Gln Ile Asp His
 20 25 30

Ala Ile Asp Ile Ile Cys Gly Phe Leu Lys Glu Arg Cys Phe Arg Gly
 35 40 45

Ser Ser Tyr Pro Val Cys Val Ser Lys Val Val Lys Gly Gly Ser Ser
 50 55 60

7

Gly Lys Gly Thr Thr Leu Arg Gly Arg Ser Asp Ala Asp Leu Val Val
 65 70 75 80

Phe Leu Ser Pro Leu Thr Thr Phe Gln Asp Gln Leu Asn Arg Arg Gly
 85 90 95

Glu Phe Ile Gln Glu Ile Arg Arg Gln Leu Glu Ala Cys Gln Arg Glu
 100 105 110

Arg Ala Leu Ser Val Lys Phe Glu Val Gln Ala Pro Arg Trp Gly Asn
 115 120 125

Pro Arg Ala Leu Ser Phe Val Leu Ser Ser Leu Gln Leu Gly Glu Gly
 130 135 140

Val Glu Phe Asp Val Leu Pro Ala Phe Asp Ala Leu Gly Gln Leu Thr
 145 150 155 160

Gly Ser Tyr Lys Pro Asn Pro Gln Ile Tyr Val Lys Leu Ile Glu Glu
 165 170 175

Cys Thr Asp Leu Gln Lys Glu Gly Glu Phe Ser Thr Cys Phe Thr Glu
 180 185 190

Leu Gln Arg Asp Phe Leu Lys Gln Arg Pro Thr Lys Leu Lys Ser Leu
 195 200 205

Ile Arg Leu Val Lys His Trp Tyr Gln Asn Cys Lys Lys Lys Leu Gly
 210 215 220

Lys Leu Pro Pro Gln Tyr Ala Leu Glu Leu Leu Thr Val Tyr Ala Trp
 225 230 235 240

Glu Arg Gly Ser Met Lys Thr His Phe Asn Thr Ala Gln Gly Phe Arg
 245 250 255

Thr Val Leu Glu Leu Val Ile Asn Tyr Gln Gln Leu Cys Ile Tyr Trp
 260 265 270

Thr Lys Tyr Tyr Asp Phe Lys Asn Pro Ile Ile Glu Lys Tyr Leu Arg
 275 280 285

Arg Gln Leu Thr Lys Pro Arg Pro Val Ile Leu Asp Pro Ala Asp Pro
 290 295 300

Thr Gly Asn Leu Gly Gly Gly Asp Pro Lys Gly Trp Arg Gln Leu Ala
 305 310 315 320

Gln Glu Ala Glu Ala Trp Leu Asn Tyr Pro Cys Phe Lys Asn Trp Asp
 325 330 335

Gly Ser Pro Val Ser Ser Trp Ile Leu Leu Ala Glu Ser Asn Ser Thr
 340 345 350

Asp Asp Glu Thr Asp Asp Pro Arg Thr Tyr Gln Lys Tyr Gly Tyr Ile
 355 360 365

Gly Thr His Glu Tyr Pro His Phe Ser His Arg Pro Ser Thr Leu Gln
 370 375 380

Ala Ala Ser Thr Pro Gln Ala Glu Glu Asp Trp Thr Cys Thr Ile Leu
 385 390 395 400

<210> 3
 <211> 1207
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(1206)

<300>
 <301> Benech et al.
 <302> Structure of two forms of the interferon-induced (2'-5') oligo A
 synthetase human cells based on cDNAs and gene sequences
 <303> EMBO J.
 <304> 4
 <305> 9
 <306> 2249-2256
 <307> 1985
 <308> NM_016816
 <309> 2003-04-06

<400> 3
 atg atg gat ctc aga aat acc cca gcc aaa tct ctg gac aag ttc att 48
 Met Met Asp Leu Arg Asn Thr Pro Ala Lys Ser Leu Asp Lys Phe Ile
 1 5 10 15
 gaa gac tat ctc ttg cca gac acg tgt ttc cgc atg caa atc gac cat 96
 Glu Asp Tyr Leu Leu Pro Asp Thr Cys Phe Arg Met Gln Ile Asp His
 20 25 30
 gcc att gac atc atc tgt ggg ttc ctg aag gaa agg tgc ttc cga ggt 144
 Ala Ile Asp Ile Ile Cys Gly Phe Leu Lys Glu Arg Cys Phe Arg Gly
 35 40 45

agc tcc tac cct gtg tgt gtg tcc aag gtg gta aag ggt ggc tcc tca Ser Ser Tyr Pro Val Cys Val Ser Lys Val Val Lys Gly Gly Ser Ser 50 55 60	192
ggc aag ggc acc acc ctc aga ggc cga tct gac gct gac ctg gtt gtc Gly Lys Gly Thr Thr Leu Arg Gly Arg Ser Asp Ala Asp Leu Val Val 65 70 75 80	240
ttc ctc agt cct ctc acc act ttt cag gat cag tta aat cgc cgg gga Phe Leu Ser Pro Leu Thr Thr Phe Gln Asp Gln Leu Asn Arg Arg Gly 85 90 95	288
gag ttc atc cag gaa att agg aga cag ctg gaa gcc tgt caa aga gag Glu Phe Ile Gln Glu Ile Arg Arg Gln Leu Glu Ala Cys Gln Arg Glu 100 105 110	336
aga gca ctt tcc gtg aag ttt gag gtc cag gct cca cgc tgg ggc aac Arg Ala Leu Ser Val Lys Phe Glu Val Gln Ala Pro Arg Trp Gly Asn 115 120 125	384
ccc cgt gcg ctc agc ttc gta ctg agt tcg ctc cag ctc ggg gag ggg Pro Arg Ala Leu Ser Phe Val Leu Ser Ser Leu Gln Leu Gly Glu Gly 130 135 140	432
gtg gag ttc gat gtg ctg cct gcc ttt gat gcc ctg ggt cag ttg act Val Glu Phe Asp Val Leu Pro Ala Phe Asp Ala Leu Gly Gln Leu Thr 145 150 155 160	480
ggc agc tat aaa cct aac ccc caa atc tat gtc aag ctc atc gag gag Gly Ser Tyr Lys Pro Asn Pro Gln Ile Tyr Val Lys Leu Ile Glu Glu 165 170 175	528
tgc acc gac ctg cag aaa gag ggc gag ttc tcc acc tgc ttc aca gaa Cys Thr Asp Leu Gln Lys Glu Gly Glu Phe Ser Thr Cys Phe Thr Glu 180 185 190	576
cta cag aga gac ttc ctg aag cag cgc ccc acc aag ctc aag agc ctc Leu Gln Arg Asp Phe Leu Lys Gln Arg Pro Thr Lys Leu Lys Ser Leu 195 200 205	624
atc cgc cta gtc aag cac tgg tac caa aat tgt aag aag aag ctt ggg Ile Arg Leu Val Lys His Trp Tyr Gln Asn Cys Lys Lys Lys Leu Gly 210 215 220	672
aag ctg cca cct cag tat gcc ctg gag ctc ctg acg gtc tat gct tgg Lys Leu Pro Pro Gln Tyr Ala Leu Glu Leu Leu Thr Val Tyr Ala Trp 225 230 235 240	720
gag cga ggg agc atg aaa aca cat ttc aac aca gcc caa gga ttt cgg Glu Arg Gly Ser Met Lys Thr His Phe Asn Thr Ala Gln Gly Phe Arg 245 250 255	768
acg gtc ttg gaa tta gtc ata aac tac cag caa ctc tgc atc tac tgg Thr Val Leu Glu Leu Val Ile Asn Tyr Gln Gln Leu Cys Ile Tyr Trp 260 265 270	816
aca aag tat tat gac ttt aaa aac ccc att att gaa aag tac ctg aga Thr Lys Tyr Tyr Asp Phe Lys Asn Pro Ile Ile Glu Lys Tyr Leu Arg	864

10

275	280	285	
agg cag ctc acg aaa ccc	agg cct gtg atc ctg gac ccg gcg gac cct		912
Arg Gln Leu Thr Lys Pro	Arg Pro Val Ile Leu Asp Pro Ala Asp Pro		
290	295	300	
aca gga aac ttg ggt ggt gga gac cca aag ggt tgg agg cag ctg gca			960
Thr Gly Asn Leu Gly Gly Gly Asp Pro Lys Gly Trp Arg Gln Leu Ala			
305	310	315	320
caa gag gct gag gcc tgg ctg aat tac cca tgc ttt aag aat tgg gat			1008
Gln Glu Ala Glu Ala Trp Leu Asn Tyr Pro Cys Phe Lys Asn Trp Asp			
	325	330	335
ggg tcc cca gtg agc tcc tgg att ctg ctg gct gaa agc aac agt aca			1056
Gly Ser Pro Val Ser Ser Trp Ile Leu Leu Ala Glu Ser Asn Ser Thr			
	340	345	350
gac gat gag acc gac gat ccc agg acg tat cag aaa tat ggt tac att			1104
Asp Asp Glu Thr Asp Asp Pro Arg Thr Tyr Gln Lys Tyr Gly Tyr Ile			
	355	360	365
gga aca cat gag tac cct cat ttc tct cat aga ccc agc acg ctc cag			1152
Gly Thr His Glu Tyr Pro His Phe Ser His Arg Pro Ser Thr Leu Gln			
	370	375	380
gca gca tcc acc cca cag gca gaa gag gac tgg acc tgc acc atc ctc			1200
Ala Ala Ser Thr Pro Gln Ala Glu Glu Asp Trp Thr Cys Thr Ile Leu			
	385	390	400
tga atg c			1207
Met			

<210> 4
 <211> 400
 <212> PRT
 <213> Homo sapiens

<400> 4

Met Met Asp Leu Arg Asn Thr Pro Ala Lys Ser Leu Asp Lys Phe Ile
1 5 10 15

Glu Asp Tyr Leu Leu Pro Asp Thr Cys Phe Arg Met Gln Ile Asp His
20 25 30

Ala Ile Asp Ile Ile Cys Gly Phe Leu Lys Glu Arg Cys Phe Arg Gly
35 40 45

Ser Ser Tyr Pro Val Cys Val Ser Lys Val Val Lys Gly Gly Ser Ser
50 55 60

Gly Lys Gly Thr Thr Leu Arg Gly Arg Ser Asp Ala Asp Leu Val Val

11

65		70		75		80									
Phe	Leu	Ser	Pro	Leu	Thr	Thr	Phe	Gln	Asp	Gln	Leu	Asn	Arg	Arg	Gly
			85						90					95	
Glu	Phe	Ile	Gln	Glu	Ile	Arg	Arg	Gln	Leu	Glu	Ala	Cys	Gln	Arg	Glu
			100					105					110		
Arg	Ala	Leu	Ser	Val	Lys	Phe	Glu	Val	Gln	Ala	Pro	Arg	Trp	Gly	Asn
		115					120					125			
Pro	Arg	Ala	Leu	Ser	Phe	Val	Leu	Ser	Ser	Leu	Gln	Leu	Gly	Glu	Gly
	130					135					140				
Val	Glu	Phe	Asp	Val	Leu	Pro	Ala	Phe	Asp	Ala	Leu	Gly	Gln	Leu	Thr
145					150					155					160
Gly	Ser	Tyr	Lys	Pro	Asn	Pro	Gln	Ile	Tyr	Val	Lys	Leu	Ile	Glu	Glu
				165					170					175	
Cys	Thr	Asp	Leu	Gln	Lys	Glu	Gly	Glu	Phe	Ser	Thr	Cys	Phe	Thr	Glu
			180					185					190		
Leu	Gln	Arg	Asp	Phe	Leu	Lys	Gln	Arg	Pro	Thr	Lys	Leu	Lys	Ser	Leu
		195					200					205			
Ile	Arg	Leu	Val	Lys	His	Trp	Tyr	Gln	Asn	Cys	Lys	Lys	Lys	Leu	Gly
	210					215					220				
Lys	Leu	Pro	Pro	Gln	Tyr	Ala	Leu	Glu	Leu	Leu	Thr	Val	Tyr	Ala	Trp
225					230					235					240
Glu	Arg	Gly	Ser	Met	Lys	Thr	His	Phe	Asn	Thr	Ala	Gln	Gly	Phe	Arg
				245					250					255	
Thr	Val	Leu	Glu	Leu	Val	Ile	Asn	Tyr	Gln	Gln	Leu	Cys	Ile	Tyr	Trp
			260					265					270		
Thr	Lys	Tyr	Tyr	Asp	Phe	Lys	Asn	Pro	Ile	Ile	Glu	Lys	Tyr	Leu	Arg
		275					280					285			
Arg	Gln	Leu	Thr	Lys	Pro	Arg	Pro	Val	Ile	Leu	Asp	Pro	Ala	Asp	Pro
	290					295					300				

12

Thr Gly Asn Leu Gly Gly Gly Asp Pro Lys Gly Trp Arg Gln Leu Ala
305 310 315 320

Gln Glu Ala Glu Ala Trp Leu Asn Tyr Pro Cys Phe Lys Asn Trp Asp
325 330 335

Gly Ser Pro Val Ser Ser Trp Ile Leu Leu Ala Glu Ser Asn Ser Thr
340 345 350

Asp Asp Glu Thr Asp Asp Pro Arg Thr Tyr Gln Lys Tyr Gly Tyr Ile
355 360 365

Gly Thr His Glu Tyr Pro His Phe Ser His Arg Pro Ser Thr Leu Gln
370 375 380

Ala Ala Ser Thr Pro Gln Ala Glu Glu Asp Trp Thr Cys Thr Ile Leu
385 390 395 400

<210> 5
<211> 2064
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (1)..(2064)

<300>
<301> Hovnanian, A., et al.
<302> The human 2', 5'-oligoadenylate synthetase locus is composed of
three distinct genes
<303> Genomics
<304> 52
<305> 3
<306> 267-277
<307> 1998
<308> NCBI/NM_002535
<309> 2003-04-03

<300>
<301> Marie, I. and Hovanessian, A.G.
<302> The 69-kDa 2-5A synthetase is composed of two homologous and
adjacent functional domains
<303> J. Biol. Chem.
<304> 267
<305> 14
<306> 9933-9939
<307> 1992
<308> NCBI/NM_002535
<309> 2003-04-03

<300>

<301> Marie, I., et al.
 <302> Differential expression and distinct structure of 69- and 100-kDa forms of 2-5A synthetase
 <303> J. Biol. Chem.
 <304> 265
 <305> 30
 <306> 18601-18607
 <307> 1990
 <308> NCBI/NM_002535
 <309> 2003-04-03

<300>
 <301> Marie, I., et al.
 <302> Preparation and characterization of polyclonal antibodies specific for the 69 and 100 k-dalton forms of human 2-5A synthetase
 <303> Biochem. Biophys. Res. Commun.
 <304> 160
 <305> 2
 <306> 580-587
 <307> 1989
 <308> NCBI/NM_002535
 <309> 2003-04-03

<300>
 <301> Hovanessian, A.G., et al.
 <302> Characterization of 69- and 100-kDa forms of 2-5A-synthetase from interferon-treated human cells
 <303> J. Biol. Chem.
 <304> 263
 <305> 10
 <306> 4959
 <307> 1988
 <308> NCBI/NM_002535
 <309> 2003-04-03

<300>
 <301> Hovanessian, A.G., et al.
 <302> Identification of 69-kd and 100-kd forms of 2-5A synthetase
 <303> EMBO J.
 <304> 6
 <305> 5
 <306> 1273-1280
 <307> 1987
 <308> NCBI/NM_002535
 <309> 2003-04-03

<400> 5
 atg gga aat ggg gag tcc cag ctg tcc tcg gtg cct gct cag aag ctg 48
 Met Gly Asn Gly Glu Ser Gln Leu Ser Ser Val Pro Ala Gln Lys Leu
 1 5 10 15
 ggt tgg ttt atc cag gaa tac ctg aag ccc tac gaa gaa tgt cag aca 96
 Gly Trp Phe Ile Gln Glu Tyr Leu Lys Pro Tyr Glu Glu Cys Gln Thr
 20 25 30
 ctg atc gac gag atg gtg aac acc atc tgt gac gtc tgc agg aac ccc 144
 Leu Ile Asp Glu Met Val Asn Thr Ile Cys Asp Val Cys Arg Asn Pro
 35 40 45

gaa cag ttc ccc ctg gtg cag gga gtg gcc ata ggt ggc tcc tat gga	192
Glu Gln Phe Pro Leu Val Gln Gly Val Ala Ile Gly Gly Ser Tyr Gly	
50 55 60	
cgg aaa aca gtc tta aga ggc aac tcc gat ggt acc ctt gtc ctt ttc	240
Arg Lys Thr Val Leu Arg Gly Asn Ser Asp Gly Thr Leu Val Leu Phe	
65 70 75 80	
ttc agt gac tta aaa caa ttc cag gat cag aag aga agc caa cgt gac	288
Phe Ser Asp Leu Lys Gln Phe Gln Asp Gln Lys Arg Ser Gln Arg Asp	
85 90 95	
atc ctc gat aaa act ggg gat aag ctg aag ttc tgt ctg ttc acg aag	336
Ile Leu Asp Lys Thr Gly Asp Lys Leu Lys Phe Cys Leu Phe Thr Lys	
100 105 110	
tgg ttg aaa aac aat ttc gag atc cag aag tcc ctt gat ggg tcc acc	384
Trp Leu Lys Asn Asn Phe Glu Ile Gln Lys Ser Leu Asp Gly Ser Thr	
115 120 125	
atc cag gtg ttc aca aaa aat cag aga atc tct ttc gag gtg ctg gcc	432
Ile Gln Val Phe Thr Lys Asn Gln Arg Ile Ser Phe Glu Val Leu Ala	
130 135 140	
gcc ttc aac gct ctg agc tta aat gat aat ccc agc ccc tgg atc tat	480
Ala Phe Asn Ala Leu Ser Leu Asn Asp Asn Pro Ser Pro Trp Ile Tyr	
145 150 155 160	
cga gag ctc aaa aga tcc ttg gat aag aca aat gcc agt cct ggt gag	528
Arg Glu Leu Lys Arg Ser Leu Asp Lys Thr Asn Ala Ser Pro Gly Glu	
165 170 175	
ttt gca gtc tgc ttc act gaa ctc cag cag aag ttt ttt gac aac cgt	576
Phe Ala Val Cys Phe Thr Glu Leu Gln Gln Lys Phe Phe Asp Asn Arg	
180 185 190	
cct gga aaa cta aag gat ttg atc ctc ttg ata aag cac tgg cat caa	624
Pro Gly Lys Leu Lys Asp Leu Ile Leu Leu Ile Lys His Trp His Gln	
195 200 205	
cag tgc cag aaa aaa atc aag gat tta ccc tcg ctg tct ccg tat gcc	672
Gln Cys Gln Lys Lys Ile Lys Asp Leu Pro Ser Leu Ser Pro Tyr Ala	
210 215 220	
ctg gag ctg ctt acg gtg tat gcc tgg gaa cag ggg tgc aga aaa gac	720
Leu Glu Leu Leu Thr Val Tyr Ala Trp Glu Gln Gly Cys Arg Lys Asp	
225 230 235 240	
aac ttt gac att gct gaa ggc gtc aga acg gtt ctg gag ctg atc aaa	768
Asn Phe Asp Ile Ala Glu Gly Val Arg Thr Val Leu Glu Leu Ile Lys	
245 250 255	
tgc cag gag aag ctg tgt atc tat tgg atg gtc aac tac aac ttt gaa	816
Cys Gln Glu Lys Leu Cys Ile Tyr Trp Met Val Asn Tyr Asn Phe Glu	
260 265 270	
gat gag acc atc agg aac atc ctg ctg cac cag ctc caa tca gcg agg	864
Asp Glu Thr Ile Arg Asn Ile Leu Leu His Gln Leu Gln Ser Ala Arg	

15

275	280	285	
cca gta atc ttg gat Pro Val Ile Leu Asp 290	cca gtt gac cca acc aat aat Pro Val Asp Pro Thr Asn Asn 295 300	gtg agt gga gat Val Ser Gly Asp	912
aaa ata tgc tgg caa tgg ctg Lys Ile Cys Trp Gln Trp Leu 305 310	aaa gaa gct caa acc tgg ttg act Lys Lys Glu Ala Gln Thr Trp Leu Thr 315 320		960
tct ccc aac ctg gat aat gag tta cct Ser Pro Asn Leu Asp Asn Glu Leu Pro 325 330	gca cca tct tgg aat gtc ctg Ala Pro Ser Trp Asn Val Leu 335		1008
cct gca cca ctc ttc acg acc cca ggc Pro Ala Pro Leu Phe Thr Thr Pro 340 345	cac ctt ctg gat aag ttc atc His Leu Leu Asp Lys Phe Ile 350		1056
aag gag ttt ctc cag ccc aac aaa tgc Lys Glu Phe Leu Gln Pro Asn Lys 355 360	tgc ttc cta gag cag att gac agt Cys Phe Leu Glu Gln Ile Asp Ser 365		1104
gct gtt aac atc atc cgt aca ttc ctt Ala Val Asn Ile Ile Arg Thr Phe 370 375	aaa gaa aac tgc ttc cga caa Lys Glu Asn Cys Phe Arg Gln 380		1152
tca aca gcc aag atc cag att gtc cgg Ser Thr Ala Lys Ile Gln Ile Val 385 390	gga gga tca acc gcc aaa ggc Gly Gly Ser Thr Ala Lys Gly 395 400		1200
aca gct ctg aag act ggc tct gat gcc Thr Ala Leu Lys Thr Gly Ser Asp 405 410	gat ctc gtc gtg ttc cat aac Ala Asp Leu Val Val Phe His Asn 415		1248
tca ctt aaa agc tac acc tcc caa Ser Leu Lys Ser Tyr Thr Ser Gln 420 425	aaa aac gag cgg cac aaa atc gtc Lys Asn Glu Arg His Lys Ile Val 430		1296
aag gaa atc cat gaa cag ctg aaa gcc Lys Glu Ile His Glu Gln Leu Lys 435 440	ttt tgg agg gag aag gag gag Ala Phe Trp Arg Glu Lys Glu Glu 445		1344
gag ctt gaa gtc agc ttt gag cct ccc Glu Leu Glu Val Ser Phe Glu Pro 450 455	aag tgg aag gct ccc agg gtg Lys Trp Lys Ala Pro Arg Val 460		1392
ctg agc ttc tct ctg aaa tcc aaa gtc Leu Ser Phe Ser Leu Lys Ser Lys 465 470	ctc aac gaa agt gtc agc ttt Leu Asn Glu Ser Val Ser Phe 475 480		1440
gat gtg ctt cct gcc ttt aat gca ctg Asp Val Leu Pro Ala Phe Asn Ala 485 490	ggt cag ctg agt tct ggc tcc Gly Gln Leu Ser Ser Gly Ser 495		1488
aca ccc agc ccc gag gtt tat gca ggg Thr Pro Ser Pro Glu Val Tyr Ala 500 505	ctc att gat ctg tat aaa tcc Leu Ile Asp Leu Tyr Lys Ser 510		1536
tcg gac ctc ccg gga gga gag ttt tct acc tgt ttc aca gtc ctg cag			1584

16

Ser Asp Leu Pro Gly Gly Glu Phe Ser Thr Cys Phe Thr Val Leu Gln
 515 520 525
 cga aac ttc att cgc tcc cgg ccc acc aaa cta aag gat tta att cgc 1632
 Arg Asn Phe Ile Arg Ser Arg Pro Thr Lys Leu Lys Asp Leu Ile Arg
 530 535 540
 ctg gtg aag cac tgg tac aaa gag tgt gaa agg aaa ctg aag cca aag 1680
 Leu Val Lys His Trp Tyr Lys Glu Cys Glu Arg Lys Leu Lys Pro Lys
 545 550 555 560
 ggg tct ttg ccc cca aag tat gcc ttg gag ctg ctc acc atc tat gcc 1728
 Gly Ser Leu Pro Pro Lys Tyr Ala Leu Glu Leu Leu Thr Ile Tyr Ala
 565 570 575
 tgg gag cag ggg agt gga gtg ccg gat ttt gac act gca gaa ggt ttc 1776
 Trp Glu Gln Gly Ser Gly Val Pro Asp Phe Asp Thr Ala Glu Gly Phe
 580 585 590
 cgg aca gtc ctg gag ctg gtc aca caa tat cag cag ctc ggc atc ttc 1824
 Arg Thr Val Leu Glu Leu Val Thr Gln Tyr Gln Gln Leu Gly Ile Phe
 595 600 605
 tgg aag gtc aat tac aac ttt gaa gat gag acc gtg agg aag ttt cta 1872
 Trp Lys Val Asn Tyr Asn Phe Glu Asp Glu Thr Val Arg Lys Phe Leu
 610 615 620
 ctg agc cag ttg cag aaa acc agg cct gtg atc ttg gac cca ggc gaa 1920
 Leu Ser Gln Leu Gln Lys Thr Arg Pro Val Ile Leu Asp Pro Gly Glu
 625 630 635 640
 ccc aca ggt gac gtg ggt gga ggg gac cgt tgg tgt tgg cat ctt ctg 1968
 Pro Thr Gly Asp Val Gly Gly Gly Asp Arg Trp Cys Trp His Leu Leu
 645 650 655
 gac aaa gaa gca aag gtt agg tta tcc tct ccc tgc ttc aag gat ggg 2016
 Asp Lys Glu Ala Lys Val Arg Leu Ser Ser Pro Cys Phe Lys Asp Gly
 660 665 670
 act gga aac cca ata cca cct tgg aaa gtg ccg gta aaa gtc atc taa 2064
 Thr Gly Asn Pro Ile Pro Pro Trp Lys Val Pro Val Lys Val Ile
 675 680 685

<210> 6
 <211> 687
 <212> PRT
 <213> Homo sapiens

<400> 6

Met Gly Asn Gly Glu Ser Gln Leu Ser Ser Val Pro Ala Gln Lys Leu
 1 5 10 15

Gly Trp Phe Ile Gln Glu Tyr Leu Lys Pro Tyr Glu Glu Cys Gln Thr
 20 25 30

17

Leu Ile Asp Glu Met Val Asn Thr Ile Cys Asp Val Cys Arg Asn Pro
 35 40 45

Glu Gln Phe Pro Leu Val Gln Gly Val Ala Ile Gly Gly Ser Tyr Gly
 50 55 60

Arg Lys Thr Val Leu Arg Gly Asn Ser Asp Gly Thr Leu Val Leu Phe
 65 70 75 80

Phe Ser Asp Leu Lys Gln Phe Gln Asp Gln Lys Arg Ser Gln Arg Asp
 85 90 95

Ile Leu Asp Lys Thr Gly Asp Lys Leu Lys Phe Cys Leu Phe Thr Lys
 100 105 110

Trp Leu Lys Asn Asn Phe Glu Ile Gln Lys Ser Leu Asp Gly Ser Thr
 115 120 125

Ile Gln Val Phe Thr Lys Asn Gln Arg Ile Ser Phe Glu Val Leu Ala
 130 135 140

Ala Phe Asn Ala Leu Ser Leu Asn Asp Asn Pro Ser Pro Trp Ile Tyr
 145 150 155 160

Arg Glu Leu Lys Arg Ser Leu Asp Lys Thr Asn Ala Ser Pro Gly Glu
 165 170 175

Phe Ala Val Cys Phe Thr Glu Leu Gln Gln Lys Phe Phe Asp Asn Arg
 180 185 190

Pro Gly Lys Leu Lys Asp Leu Ile Leu Leu Ile Lys His Trp His Gln
 195 200 205

Gln Cys Gln Lys Lys Ile Lys Asp Leu Pro Ser Leu Ser Pro Tyr Ala
 210 215 220

Leu Glu Leu Leu Thr Val Tyr Ala Trp Glu Gln Gly Cys Arg Lys Asp
 225 230 235 240

Asn Phe Asp Ile Ala Glu Gly Val Arg Thr Val Leu Glu Leu Ile Lys
 245 250 255

Cys Gln Glu Lys Leu Cys Ile Tyr Trp Met Val Asn Tyr Asn Phe Glu
 260 265 270

Asp Glu Thr Ile Arg Asn Ile Leu Leu His Gln Leu Gln Ser Ala Arg
 275 280 285

Pro Val Ile Leu Asp Pro Val Asp Pro Thr Asn Asn Val Ser Gly Asp
 290 295 300

Lys Ile Cys Trp Gln Trp Leu Lys Lys Glu Ala Gln Thr Trp Leu Thr
 305 310 315 320

Ser Pro Asn Leu Asp Asn Glu Leu Pro Ala Pro Ser Trp Asn Val Leu
 325 330 335

Pro Ala Pro Leu Phe Thr Thr Pro Gly His Leu Leu Asp Lys Phe Ile
 340 345 350

Lys Glu Phe Leu Gln Pro Asn Lys Cys Phe Leu Glu Gln Ile Asp Ser
 355 360 365

Ala Val Asn Ile Ile Arg Thr Phe Leu Lys Glu Asn Cys Phe Arg Gln
 370 375 380

Ser Thr Ala Lys Ile Gln Ile Val Arg Gly Gly Ser Thr Ala Lys Gly
 385 390 395 400

Thr Ala Leu Lys Thr Gly Ser Asp Ala Asp Leu Val Val Phe His Asn
 405 410 415

Ser Leu Lys Ser Tyr Thr Ser Gln Lys Asn Glu Arg His Lys Ile Val
 420 425 430

Lys Glu Ile His Glu Gln Leu Lys Ala Phe Trp Arg Glu Lys Glu Glu
 435 440 445

Glu Leu Glu Val Ser Phe Glu Pro Pro Lys Trp Lys Ala Pro Arg Val
 450 455 460

Leu Ser Phe Ser Leu Lys Ser Lys Val Leu Asn Glu Ser Val Ser Phe
 465 470 475 480

Asp Val Leu Pro Ala Phe Asn Ala Leu Gly Gln Leu Ser Ser Gly Ser
 485 490 495

Thr Pro Ser Pro Glu Val Tyr Ala Gly Leu Ile Asp Leu Tyr Lys Ser
 500 505 510

Ser Asp Leu Pro Gly Gly Glu Phe Ser Thr Cys Phe Thr Val Leu Gln
 515 520 525

Arg Asn Phe Ile Arg Ser Arg Pro Thr Lys Leu Lys Asp Leu Ile Arg
 530 535 540

Leu Val Lys His Trp Tyr Lys Glu Cys Glu Arg Lys Leu Lys Pro Lys
 545 550 555 560

Gly Ser Leu Pro Pro Lys Tyr Ala Leu Glu Leu Leu Thr Ile Tyr Ala
 565 570 575

Trp Glu Gln Gly Ser Gly Val Pro Asp Phe Asp Thr Ala Glu Gly Phe
 580 585 590

Arg Thr Val Leu Glu Leu Val Thr Gln Tyr Gln Gln Leu Gly Ile Phe
 595 600 605

Trp Lys Val Asn Tyr Asn Phe Glu Asp Glu Thr Val Arg Lys Phe Leu
 610 615 620

Leu Ser Gln Leu Gln Lys Thr Arg Pro Val Ile Leu Asp Pro Gly Glu
 625 630 635 640

Pro Thr Gly Asp Val Gly Gly Gly Asp Arg Trp Cys Trp His Leu Leu
 645 650 655

Asp Lys Glu Ala Lys Val Arg Leu Ser Ser Pro Cys Phe Lys Asp Gly
 660 665 670

Thr Gly Asn Pro Ile Pro Pro Trp Lys Val Pro Val Lys Val Ile
 675 680 685

<210> 7

<211> 2186

<212> DNA

<213> Homo sapiens

<300>

<301> Marie, I. and Hovanessian, A.G.

<302> The 69-kDa 2-5A synthetase is composed of two homologous and adjacent functional domains

<303> J. Biol. Chem.

<304> 267

<305> 14

<306> 9933-9939

<307> 1992

<308> (unknown)

<309> 2003-04-03

<400> 7

atgggaaatg gggagtccca gctgtcctcg gtgcctgctc agaagctggg ttggtttatc	60
caggaatacc tgaagcccta cgaagaatgt cagacactga tcgacgagat ggtgaacacc	120
atctgtgacg tctgcaggaa ccccgaaacag ttccccctgg tgcaggaggat ggccataggt	180
ggctcctatg gacggaaaac agtcttaaga ggcaactccg atggtaccct tgtccttttc	240
ttcagtgact taaaacaatt ccaggatcag aagagaagcc aacgtgacat cctcgataaa	300
actggggata agctgaagtt ctgtctgttc acgaagtggg tgaaaaacaa ttctcgagatc	360
cagaagtccc ttgatgggtc caccatccag gtgttcacaa aaaatcagag aatctctttc	420
gaggtgctgg ccgccttcaa cgctctgagc ttaaatagata atcccagccc ctggatctat	480
cgagagctca aaagatcctt ggataagaca aatgccagtc ctggtgagtt tgcagtctgc	540
ttcactgaac tccagcagaa gttttttgac aaccgtcctg gaaaactaaa ggatttgatc	600
ctcttgataa agcactggca tcaacagtgc cagaaaaaaa tcaaggattt accctcgctg	660
tctccgtatg ccctggagct gcttacggtg tatgcctggg aacaggggtg cagaaaagac	720
aactttgaca ttgctgaagg cgtcagaacg gttctggagc tgatcaaata ccaggagaag	780
ctgtgtatct attggatggg caactacaac tttgaagatg agaccatcag gaacatcctg	840
ctgcaccagc tccaatcagc gaggccagta atcttgatc cagttgacct aaccaataat	900
gtgagtggag ataaaaatat ctggcaatgg ctgaaaaaag aagctcaaac ctggttgact	960
tctcccaacc tggataatga gttacctgca ccatcttgga atgtcctgcc tgcaccactc	1020
ttcacgacct caggccacct tctggataag ttcatcaagg agtttctcca gcccacaaa	1080
tgcttcctag agcagattga cagtgtgtt aacatcatcc gtacattcct taaagaaaac	1140
tgcttcogac aatcaacagc caagatccag attgtccggg gaggatcaac cgccaaaggc	1200
acagctctga agactggctc tgatgccgat ctcgctgtgt tccataactc acttaaaagc	1260
tacacctccc aaaaaaacga gcggcacaaa atcgtcaagg aaatccatga acagctgaaa	1320
gccttttgga gggagaagga ggaggagctt gaagtcagct ttgagcctcc caagtggaag	1380
gctcccaggg tgctgagctt ctctctgaaa tccaaagtcc tcaacgaaag tgtcagcttt	1440
gatgtgcttc ctgcctttaa tgcactgggt cagctgagtt ctggctccac acccagcccc	1500
gaggtttatg cagggtcat tgatctgtat aaatcctcgg acctcccggg aggagagttt	1560
tctacctgtt tcacagtcct gcagcgaaac ttcattcgct cccggccac caaactaaag	1620

21

```

gattttaattc gcctggtgaa gcactggtac aaagagtgtg aaaggaaact gaagccaaag 1680
gggtcttttgc ccccaaagta tgccttggag ctgctcacca tctatgcctg ggagcagggg 1740
agtggagtg cggattttga cactgcagaa ggtttccgga cagtccctgga gctggtcaca 1800
caatatcagc agctcggcat cttctggaag gtcaattaca actttgaaga tgagaccgtg 1860
aggaagtttc tactgagcca gttgcagaaa accaggcctg tgatcttgga ccagggcgaa 1920
cccacaggtg acgtgggtgg aggggaccgt tgggtgttggc atcttctgga caaagaagca 1980
aaggttaggt tctcctctcc ctgcttcaag gatgggactg gaaaccaat accaccttgg 2040
aaagtgccga caatgcagac accaggaagt tgtggagcta ggattccatc ctattgtcaa 2100
tgagatgttg tcatccagaa gccatagaat cctgaataat aattctaaaa gaaacttctg 2160
gagatcatct ggcaatcgct tttaaa 2186

```

```

<210> 8
<211> 727
<212> PRT
<213> Homo sapiens

```

```

<400> 8

```

```

Met Gly Asn Gly Glu Ser Gln Leu Ser Ser Val Pro Ala Gln Lys Leu
1           5           10           15

```

```

Gly Trp Phe Ile Gln Glu Tyr Leu Lys Pro Tyr Glu Glu Cys Gln Thr
          20           25           30

```

```

Leu Ile Asp Glu Met Val Asn Thr Ile Cys Asp Val Cys Arg Asn Pro
          35           40           45

```

```

Glu Gln Phe Pro Leu Val Gln Gly Val Ala Ile Gly Gly Ser Tyr Gly
          50           55           60

```

```

Arg Lys Thr Val Leu Arg Gly Asn Ser Asp Gly Thr Leu Val Leu Phe
65           70           75           80

```

```

Phe Ser Asp Leu Lys Gln Phe Gln Asp Gln Lys Arg Ser Gln Arg Asp
          85           90           95

```

```

Ile Leu Asp Lys Thr Gly Asp Lys Leu Lys Phe Cys Leu Phe Thr Lys
          100          105          110

```

```

Trp Leu Lys Asn Asn Phe Glu Ile Gln Lys Ser Leu Asp Gly Ser Thr
          115          120          125

```


Ile Gln Val Phe Thr Lys Asn Gln Arg Ile Ser Phe Glu Val Leu Ala
 130 135 140

Ala Phe Asn Ala Leu Ser Leu Asn Asp Asn Pro Ser Pro Trp Ile Tyr
 145 150 155 160

Arg Glu Leu Lys Arg Ser Leu Asp Lys Thr Asn Ala Ser Pro Gly Glu
 165 170 175

Phe Ala Val Cys Phe Thr Glu Leu Gln Gln Lys Phe Phe Asp Asn Arg
 180 185 190

Pro Gly Lys Leu Lys Asp Leu Ile Leu Leu Ile Lys His Trp His Gln
 195 200 205

Gln Cys Gln Lys Lys Ile Lys Asp Leu Pro Ser Leu Ser Pro Tyr Ala
 210 215 220

Leu Glu Leu Leu Thr Val Tyr Ala Trp Glu Gln Gly Cys Arg Lys Asp
 225 230 235 240

Asn Phe Asp Ile Ala Glu Gly Val Arg Thr Val Leu Glu Leu Ile Lys
 245 250 255

Cys Gln Glu Lys Leu Cys Ile Tyr Trp Met Val Asn Tyr Asn Phe Glu
 260 265 270

Asp Glu Thr Ile Arg Asn Ile Leu Leu His Gln Leu Gln Ser Ala Arg
 275 280 285

Pro Val Ile Leu Asp Pro Val Asp Pro Thr Asn Asn Val Ser Gly Asp
 290 295 300

Lys Ile Cys Trp Gln Trp Leu Lys Lys Glu Ala Gln Thr Trp Leu Thr
 305 310 315 320

Ser Pro Asn Leu Asp Asn Glu Leu Pro Ala Pro Ser Trp Asn Val Leu
 325 330 335

Pro Ala Pro Leu Phe Thr Thr Pro Gly His Leu Leu Asp Lys Phe Ile
 340 345 350

Lys Glu Phe Leu Gln Pro Asn Lys Cys Phe Leu Glu Gln Ile Asp Ser
 355 360 365

Ala Val Asn Ile Ile Arg Thr Phe Leu Lys Glu Asn Cys Phe Arg Gln
 370 375 380

Ser Thr Ala Lys Ile Gln Ile Val Arg Gly Gly Ser Thr Ala Lys Gly
 385 390 395 400

Thr Ala Leu Lys Thr Gly Ser Asp Ala Asp Leu Val Val Phe His Asn
 405 410 415

Ser Leu Lys Ser Tyr Thr Ser Gln Lys Asn Glu Arg His Lys Ile Val
 420 425 430

Lys Glu Ile His Glu Gln Leu Lys Ala Phe Trp Arg Glu Lys Glu Glu
 435 440 445

Glu Leu Glu Val Ser Phe Glu Pro Pro Lys Trp Lys Ala Pro Arg Val
 450 455 460

Leu Ser Phe Ser Leu Lys Ser Lys Val Leu Asn Glu Ser Val Ser Phe
 465 470 475 480

Asp Val Leu Pro Ala Phe Asn Ala Leu Gly Gln Leu Ser Ser Gly Ser
 485 490 495

Thr Pro Ser Pro Glu Val Tyr Ala Gly Leu Ile Asp Leu Tyr Lys Ser
 500 505 510

Ser Asp Leu Pro Gly Gly Glu Phe Ser Thr Cys Phe Thr Val Leu Gln
 515 520 525

Arg Asn Phe Ile Arg Ser Arg Pro Thr Lys Leu Lys Asp Leu Ile Arg
 530 535 540

Leu Val Lys His Trp Tyr Lys Glu Cys Glu Arg Lys Leu Lys Pro Lys
 545 550 555 560

Gly Ser Leu Pro Pro Lys Tyr Ala Leu Glu Leu Leu Thr Ile Tyr Ala
 565 570 575

Trp Glu Gln Gly Ser Gly Val Pro Asp Phe Asp Thr Ala Glu Gly Phe
 580 585 590

Arg Thr Val Leu Glu Leu Val Thr Gln Tyr Gln Gln Leu Gly Ile Phe

24

595		600		605											
Trp	Lys	Val	Asn	Tyr	Asn	Phe	Glu	Asp	Glu	Thr	Val	Arg	Lys	Phe	Leu
610						615					620				
Leu	Ser	Gln	Leu	Gln	Lys	Thr	Arg	Pro	Val	Ile	Leu	Asp	Pro	Gly	Glu
625					630					635					640
Pro	Thr	Gly	Asp	Val	Gly	Gly	Gly	Asp	Arg	Trp	Cys	Trp	His	Leu	Leu
				645					650					655	
Asp	Lys	Glu	Ala	Lys	Val	Arg	Leu	Ser	Ser	Pro	Cys	Phe	Lys	Asp	Gly
			660					665						670	
Thr	Gly	Asn	Pro	Ile	Pro	Pro	Trp	Lys	Val	Pro	Thr	Met	Gln	Thr	Pro
		675					680					685			
Gly	Ser	Cys	Gly	Ala	Arg	Ile	His	Pro	Ile	Val	Asn	Glu	Met	Phe	Ser
	690					695					700				
Ser	Arg	Ser	His	Arg	Ile	Leu	Asn	Asn	Asn	Ser	Lys	Arg	Asn	Phe	Thr
705					710					715					720
Arg	Ser	Ser	Gly	Asn	Arg	Phe									
				725											

<210> 9
 <211> 3264
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(3264)

<300>
 <301> Rebouillat, D., et al.
 <302> The 100-kDa 2',5'-oligoadenylate synthetase catalyzing preferentially the synthesis of dimeric pppA2'p5'A molecules
 <303> J. Biol. Chem.
 <304> 274
 <305> 3
 <306> 1557-1565
 <307> 1999
 <308> NCBI/AF_063613
 <309> 1999-05-04

<300>
 <301> Rebouillat, D. and Hovanessian, A.G.

25

```

<302> Direct Submission
<303> Submitted (07-May-1998) Dept.of Aids and Retroviruses, Institut
      Pasteur
<304> 0
<305> 0
<306> 0
<307> 1998
<308> NCBI/AF_063613
<309> 1999-05-04

<400> 9
atg gac ttg tac agc acc ccg gcc gct gcg ctg gac agg ttc gtg gcc      48
Met Asp Leu Tyr Ser Thr Pro Ala Ala Ala Leu Asp Arg Phe Val Ala
1              5              10              15

aga agg ctg cag ccg cgg aag gag ttc gta gag aag gcg cgg cgc gct      96
Arg Arg Leu Gln Pro Arg Lys Glu Phe Val Glu Lys Ala Arg Arg Ala
              20              25              30

ctg ggc gcc ctg gcc gct gcc ctg agg gag cgc ggg ggc cgc ctc ggt      144
Leu Gly Ala Leu Ala Ala Ala Leu Arg Glu Arg Gly Gly Arg Leu Gly
              35              40              45

gct gct gcc ccg cgg gtg ctg aaa act gtc aag gga ggc tcc tcg ggc      192
Ala Ala Ala Pro Arg Val Leu Lys Thr Val Lys Gly Gly Ser Ser Gly
              50              55              60

cgg ggc aca gct ctc aag ggt ggc tgt gat tct gaa ctt gtc atc ttc      240
Arg Gly Thr Ala Leu Lys Gly Gly Cys Asp Ser Glu Leu Val Ile Phe
65              70              75              80

ctc gac tgc ttc aag agc tat gtg gac cag agg gcc cgc cgt gca gag      288
Leu Asp Cys Phe Lys Ser Tyr Val Asp Gln Arg Ala Arg Arg Ala Glu
              85              90              95

atc ctc agt gag atg ccg gca tcg ctg gaa tcc tgg tgg cag aac cca      336
Ile Leu Ser Glu Met Arg Ala Ser Leu Glu Ser Trp Trp Gln Asn Pro
              100              105              110

gtc cct ggt ctg aga ctc acg ttt cct gag cag agc gtg cct ggg gcc      384
Val Pro Gly Leu Arg Leu Thr Phe Pro Glu Gln Ser Val Pro Gly Ala
              115              120              125

ctg cag ttc cgc ctg aca tcc gta gat ctt gag gac tgg atg gat gtt      432
Leu Gln Phe Arg Leu Thr Ser Val Asp Leu Glu Asp Trp Met Asp Val
              130              135              140

agc ctg gtg cct gcc ttc aat gtc ctg ggt cag gcc ggc tcc gcg gtc      480
Ser Leu Val Pro Ala Phe Asn Val Leu Gly Gln Ala Gly Ser Ala Val
145              150              155              160

aaa ccc aag cca caa gtc tac tct acc ctc ctc aac agt ggc tgc caa      528
Lys Pro Lys Pro Gln Val Tyr Ser Thr Leu Leu Asn Ser Gly Cys Gln
              165              170              175

ggg ggc gag cat gcg gcc tgc ttc aca gag ctg cgg agg aac ttt gtg      576
Gly Gly Glu His Ala Ala Cys Phe Thr Glu Leu Arg Arg Asn Phe Val
              180              185              190

```

26

aac att cgc cca gcc aag ttg aag aac cta atc ttg ctg gtg aag cac	624
Asn Ile Arg Pro Ala Lys Leu Lys Asn Leu Ile Leu Leu Val Lys His	
195 200 205	
tgg tac cac cag gtg tgc cta cag ggg ttg tgg aag gag acg ctg ccc	672
Trp Tyr His Gln Val Cys Leu Gln Gly Leu Trp Lys Glu Thr Leu Pro	
210 215 220	
ccg gtc tat gcc ctg gaa ttg ctg acc atc ttc gcc tgg gag cag ggc	720
Pro Val Tyr Ala Leu Glu Leu Leu Thr Ile Phe Ala Trp Glu Gln Gly	
225 230 235 240	
tgt aag aag gat gct ttc agc cta ggc gaa ggc ctc cga act gtc ctg	768
Cys Lys Lys Asp Ala Phe Ser Leu Gly Glu Gly Leu Arg Thr Val Leu	
245 250 255	
ggc ctg atc caa cag cat cag cac ctg tgt gtt ttc tgg act gtc aac	816
Gly Leu Ile Gln Gln His Gln His Leu Cys Val Phe Trp Thr Val Asn	
260 265 270	
tat ggc ttc gag gac cct gca gtt ggg cag ttc ttg cag cgg cac gtt	864
Tyr Gly Phe Glu Asp Pro Ala Val Gly Gln Phe Leu Gln Arg His Val	
275 280 285	
aag aga ccc agg cct gtg atc ctg gac cca gct gac ccc aca tgg gac	912
Lys Arg Pro Arg Pro Val Ile Leu Asp Pro Ala Asp Pro Thr Trp Asp	
290 295 300	
ctg ggg aat ggg gca gcc tgg cac tgg gat ttg cat gcc cag gag gca	960
Leu Gly Asn Gly Ala Ala Trp His Trp Asp Leu His Ala Gln Glu Ala	
305 310 315 320	
gca tcc tgc tat gac cac cca tgc ttt ctg agg ggg atg ggg gac cca	1008
Ala Ser Cys Tyr Asp His Pro Cys Phe Leu Arg Gly Met Gly Asp Pro	
325 330 335	
gtg cag tct tgg aag ggg ccg ggc ctt cca cgt gct gga tgc tca ggt	1056
Val Gln Ser Trp Lys Gly Pro Gly Leu Pro Arg Ala Gly Cys Ser Gly	
340 345 350	
ttg ggc cac ccc atc cag cta gac cct aac cag aag acc cct gaa aac	1104
Leu Gly His Pro Ile Gln Leu Asp Pro Asn Gln Lys Thr Pro Glu Asn	
355 360 365	
agc aag agc ctc aat gct gtg tac cca aga gca ggg agc aaa cct ccc	1152
Ser Lys Ser Leu Asn Ala Val Tyr Pro Arg Ala Gly Ser Lys Pro Pro	
370 375 380	
tca tgc cca gct cct ggc ccc act gcg gag cca gca tcg tac ccc tct	1200
Ser Cys Pro Ala Pro Gly Pro Thr Ala Glu Pro Ala Ser Tyr Pro Ser	
385 390 395 400	
gtg ccg gga atg gcc ttg gac ctg tct cag atc ccc acc aag gag ctg	1248
Val Pro Gly Met Ala Leu Asp Leu Ser Gln Ile Pro Thr Lys Glu Leu	
405 410 415	
gac cgc ttc atc cag gac cac ctg aag ccg agc ccc cag ttc cag gag	1296
Asp Arg Phe Ile Gln Asp His Leu Lys Pro Ser Pro Gln Phe Gln Glu	
420 425 430	

cag	gtg	aaa	aag	gcc	atc	gac	atc	atc	ttg	cgc	tgc	ctc	cat	gag	aac	1344
Gln	Val	Lys	Lys	Ala	Ile	Asp	Ile	Ile	Leu	Arg	Cys	Leu	His	Glu	Asn	
		435					440					445				
tgt	gtt	cac	aag	gcc	tca	aga	gtc	agt	aaa	ggg	ggc	tca	ttt	ggc	cgg	1392
Cys	Val	His	Lys	Ala	Ser	Arg	Val	Ser	Lys	Gly	Gly	Ser	Phe	Gly	Arg	
	450					455					460					
ggc	aca	gac	cta	agg	gat	ggc	tgt	gat	gtt	gaa	ctc	atc	atc	ttc	ctc	1440
Gly	Thr	Asp	Leu	Arg	Asp	Gly	Cys	Asp	Val	Glu	Leu	Ile	Ile	Phe	Leu	
465					470					475					480	
aac	tgc	ttc	acg	gac	tac	aag	gac	cag	ggg	ccc	cgc	cgc	gca	gag	atc	1488
Asn	Cys	Phe	Thr	Asp	Tyr	Lys	Asp	Gln	Gly	Pro	Arg	Arg	Ala	Glu	Ile	
				485					490					495		
ctt	gat	gag	atg	cga	gcg	cac	gta	gaa	tcc	tgg	tgg	cag	gac	cag	gtg	1536
Leu	Asp	Glu	Met	Arg	Ala	His	Val	Glu	Ser	Trp	Trp	Gln	Asp	Gln	Val	
			500					505					510			
ccc	agc	ctg	agc	ctt	cag	ttt	cct	gag	cag	aat	gtg	cct	gag	gct	ctg	1584
Pro	Ser	Leu	Ser	Leu	Gln	Phe	Pro	Glu	Gln	Asn	Val	Pro	Glu	Ala	Leu	
		515					520					525				
cag	ttc	cag	ctg	gtg	tcc	aca	gcc	ctg	aag	agc	tgg	acg	gat	gtt	agc	1632
Gln	Phe	Gln	Leu	Val	Ser	Thr	Ala	Leu	Lys	Ser	Trp	Thr	Asp	Val	Ser	
	530					535					540					
ctg	ctg	cct	gcc	ttc	gat	gct	gtg	ggg	cag	ctc	agt	tct	ggc	acc	aaa	1680
Leu	Leu	Pro	Ala	Phe	Asp	Ala	Val	Gly	Gln	Leu	Ser	Ser	Gly	Thr	Lys	
545					550					555					560	
cca	aat	ccc	cag	gtc	tac	tcg	agg	ctc	ctc	acc	agt	ggc	tgc	cag	gag	1728
Pro	Asn	Pro	Gln	Val	Tyr	Ser	Arg	Leu	Leu	Thr	Ser	Gly	Cys	Gln	Glu	
				565					570					575		
ggc	gag	cat	aag	gcc	tgc	ttc	gca	gag	ctg	cgg	agg	aac	ttc	atg	aac	1776
Gly	Glu	His	Lys	Ala	Cys	Phe	Ala	Glu	Leu	Arg	Arg	Asn	Phe	Met	Asn	
			580					585					590			
att	cgc	cct	gtc	aag	ctg	aag	aac	ctg	att	ctg	ctg	gtg	aag	cac	tgg	1824
Ile	Arg	Pro	Val	Lys	Leu	Lys	Asn	Leu	Ile	Leu	Leu	Val	Lys	His	Trp	
		595					600					605				
tac	cgc	cag	gtt	gcg	gct	cag	aac	aaa	gga	aaa	gga	cca	gcc	cct	gcc	1872
Tyr	Arg	Gln	Val	Ala	Ala	Gln	Asn	Lys	Gly	Lys	Gly	Pro	Ala	Pro	Ala	
	610					615					620					
tct	ctg	ccc	cca	gcc	tat	gcc	ctg	gag	ctc	ctc	acc	atc	ttt	gcc	tgg	1920
Ser	Leu	Pro	Pro	Ala	Tyr	Ala	Leu	Glu	Leu		Thr	Ile	Phe	Ala	Trp	
625					630					635					640	
gag	cag	ggc	tgc	agg	cag	gat	tgt	ttc	aac	atg	gcc	caa	ggc	ttc	cgg	1968
Glu	Gln	Gly	Cys	Arg	Gln	Asp	Cys	Phe	Asn	Met	Ala	Gln	Gly	Phe	Arg	
				645					650					655		
acg	gtg	ctg	ggg	ctc	gtg	caa	cag	cat	cag	cag	ctc	tgt	gtc	tac	tgg	2016
Thr	Val	Leu	Gly	Leu	Val	Gln	Gln	His	Gln	Gln	Leu	Cys	Val	Tyr	Trp	

660						665						670						
acg	gtc	aac	tat	agc	act	gag	gac	cca	gcc	atg	aga	atg	cac	ctt	ctt			2064
Thr	Val	Asn	Tyr	Ser	Thr	Glu	Asp	Pro	Ala	Met	Arg	Met	His	Leu	Leu			
		675					680					685						
ggc	cag	ctt	cga	aaa	ccc	aga	ccc	ctg	gtc	ctg	gac	ccc	gct	gat	ccc			2112
Gly	Gln	Leu	Arg	Lys	Pro	Arg	Pro	Leu	Val	Leu	Asp	Pro	Ala	Asp	Pro			
	690					695					700							
acc	tgg	aac	gtg	ggc	cac	ggc	agc	tgg	gag	ctg	ttg	gcc	cag	gaa	gca			2160
Thr	Trp	Asn	Val	Gly	His	Gly	Ser	Trp	Glu	Leu	Leu	Ala	Gln	Glu	Ala			
705					710				715					720				
gca	gcg	ctg	ggg	atg	cag	gcc	tgc	ttt	ctg	agt	aga	gac	ggg	aca	tct			2208
Ala	Ala	Leu	Gly	Met	Gln	Ala	Cys	Phe	Leu	Ser	Arg	Asp	Gly	Thr	Ser			
				725					730					735				
gtg	cag	ccc	tgg	gat	gtg	atg	cca	gcc	ctc	ctt	tac	caa	acc	cca	gct			2256
Val	Gln	Pro	Trp	Asp	Val	Met	Pro	Ala	Leu	Leu	Tyr	Gln	Thr	Pro	Ala			
			740					745					750					
ggg	gac	ctt	gac	aag	ttc	atc	agt	gaa	ttt	ctc	cag	ccc	aac	cgc	cag			2304
Gly	Asp	Leu	Asp	Lys	Phe	Ile	Ser	Glu	Phe	Leu	Gln	Pro	Asn	Arg	Gln			
	755						760					765						
ttc	ctg	gcc	cag	gtg	aac	aag	gcc	gtt	gat	acc	atc	tgt	tca	ttt	ttg			2352
Phe	Leu	Ala	Gln	Val	Asn	Lys	Ala	Val	Asp	Thr	Ile	Cys	Ser	Phe	Leu			
	770					775					780							
aag	gaa	aac	tgc	ttc	cgg	aat	tct	ccc	atc	aaa	gtg	atc	aag	gtg	gtc			2400
Lys	Glu	Asn	Cys	Phe	Arg	Asn	Ser	Pro	Ile	Lys	Val	Ile	Lys	Val	Val			
785					790				795					800				
aag	ggt	ggc	tct	tca	gcc	aaa	ggc	aca	gct	ctg	cga	ggc	cgc	tca	gat			2448
Lys	Gly	Gly	Ser	Ser	Ala	Lys	Gly	Thr	Ala	Leu	Arg	Gly	Arg	Ser	Asp			
			805					810					815					
gcc	gac	ctc	gtg	gtg	ttc	ctc	agc	tgc	ttc	agc	cag	ttc	act	gag	cag			2496
Ala	Asp	Leu	Val	Val	Phe	Leu	Ser	Cys	Phe	Ser	Gln	Phe	Thr	Glu	Gln			
			820					825					830					
ggc	aac	aag	cgg	gcc	gag	atc	atc	tcc	gag	atc	cga	gcc	cag	ctg	gag			2544
Gly	Asn	Lys	Arg	Ala	Glu	Ile	Ile	Ser	Glu	Ile	Arg	Ala	Gln	Leu	Glu			
		835					840					845						
gca	tgt	caa	cag	gag	cgg	cag	ttc	gag	gtc	aag	ttt	gaa	gtc	tcc	aaa			2592
Ala	Cys	Gln	Gln	Glu	Arg	Gln	Phe	Glu	Val	Lys	Phe	Glu	Val	Ser	Lys			
	850					855					860							
tgg	gag	aat	ccc	cgc	gtg	ctg	agc	ttc	tca	ctg	aca	tcc	cag	acg	atg			2640
Trp	Glu	Asn	Pro	Arg	Val	Leu	Ser	Phe	Ser	Leu	Thr	Ser	Gln	Thr	Met			
865					870				875					880				
ctg	gac	cag	agt	gtg	gac	ttt	gat	gtg	ctg	cca	gcc	ttt	gac	gcc	cta			2688
Leu	Asp	Gln	Ser	Val	Asp	Phe	Asp	Val	Leu	Pro	Ala	Phe	Asp	Ala	Leu			
				885				890					895					
ggc	cag	ctg	gtc	tct	ggc	tcc	agg	ccc	agc	tct	caa	gtc	tac	gtc	gac			2736

29

Gly	Gln	Leu	Val	Ser	Gly	Ser	Arg	Pro	Ser	Ser	Gln	Val	Tyr	Val	Asp	
			900					905					910			
ctc	atc	cac	agc	tac	agc	aat	gcg	ggc	gag	tac	tcc	acc	tgc	ttc	aca	2784
Leu	Ile	His	Ser	Tyr	Ser	Asn	Ala	Gly	Glu	Tyr	Ser	Thr	Cys	Phe	Thr	
		915					920					925				
gag	cta	caa	cgg	gac	ttc	atc	atc	tct	cgc	cct	acc	aag	ctg	aag	agc	2832
Glu	Leu	Gln	Arg	Asp	Phe	Ile	Ile	Ser	Arg	Pro	Thr	Lys	Leu	Lys	Ser	
	930					935					940					
ctg	atc	cgg	ctg	gtg	aag	cac	tgg	tac	cag	cag	tgt	acc	aag	atc	tcc	2880
Leu	Ile	Arg	Leu	Val	Lys	His	Trp	Tyr	Gln	Gln	Cys	Thr	Lys	Ile	Ser	
945					950				955					960		
aag	ggg	aga	ggc	tcc	cta	ccc	cca	cag	cac	ggg	ctg	gaa	ctc	ctg	act	2928
Lys	Gly	Arg	Gly	Ser	Leu	Pro	Pro	Gln	His	Gly	Leu	Glu	Leu	Leu	Thr	
			965					970					975			
gtg	tat	gcc	tgg	gag	cag	ggc	ggg	aag	gac	tcc	cag	ttc	aac	atg	gct	2976
Val	Tyr	Ala	Trp	Glu	Gln	Gly	Gly	Lys	Asp	Ser	Gln	Phe	Asn	Met	Ala	
		980					985					990				
gag	ggc	ttc	cgc	acg	gtc	ctg	gag	ctg	gtc	acc	cag	tac	cgc	cag	ctc	3024
Glu	Gly	Phe	Arg	Thr	Val	Leu	Glu	Leu	Val	Thr	Gln	Tyr	Arg	Gln	Leu	
	995					1000					1005					
tgt	atc	tac	tgg	acc	atc	aac	tac	aac	gcc	aag	gac	aag	act	gtt		3069
Cys	Ile	Tyr	Trp	Thr	Ile	Asn	Tyr	Asn	Ala	Lys	Asp	Lys	Thr	Val		
	1010					1015					1020					
gga	gac	ttc	ctg	aaa	cag	cag	ctt	cag	aag	ccc	agg	cct	atc	atc		3114
Gly	Asp	Phe	Leu	Lys	Gln	Gln	Leu	Gln	Lys	Pro	Arg	Pro	Ile	Ile		
	1025					1030					1035					
ctg	gat	ccg	gct	gac	ccg	aca	ggc	aac	ctg	ggc	cac	aat	gcc	cgc		3159
Leu	Asp	Pro	Ala	Asp	Pro	Thr	Gly	Asn	Leu	Gly	His	Asn	Ala	Arg		
	1040					1045					1050					
tgg	gac	ctg	ctg	gcc	aag	gaa	gct	gca	gcc	tgc	aca	tct	gcc	ctg		3204
Trp	Asp	Leu	Leu	Ala	Lys	Glu	Ala	Ala	Ala	Cys	Thr	Ser	Ala	Leu		
	1055					1060					1065					
tgc	tgc	atg	gga	cgg	aat	ggc	atc	ccc	atc	cag	cca	tgg	cca	gtg		3249
Cys	Cys	Met	Gly	Arg	Asn	Gly	Ile	Pro	Ile	Gln	Pro	Trp	Pro	Val		
	1070					1075					1080					
aag	gct	gct	gtg	tga												3264
Lys	Ala	Ala	Val													
	1085															

<210> 10
 <211> 1087
 <212> PRT
 <213> Homo sapiens
 <400> 10

30

Met Asp Leu Tyr Ser Thr Pro Ala Ala Ala Leu Asp Arg Phe Val Ala
 1 5 10 15
 Arg Arg Leu Gln Pro Arg Lys Glu Phe Val Glu Lys Ala Arg Arg Ala
 20 25 30
 Leu Gly Ala Leu Ala Ala Ala Leu Arg Glu Arg Gly Gly Arg Leu Gly
 35 40 45
 Ala Ala Ala Pro Arg Val Leu Lys Thr Val Lys Gly Gly Ser Ser Gly
 50 55 60
 Arg Gly Thr Ala Leu Lys Gly Gly Cys Asp Ser Glu Leu Val Ile Phe
 65 70 75 80
 Leu Asp Cys Phe Lys Ser Tyr Val Asp Gln Arg Ala Arg Arg Ala Glu
 85 90 95
 Ile Leu Ser Glu Met Arg Ala Ser Leu Glu Ser Trp Trp Gln Asn Pro
 100 105 110
 Val Pro Gly Leu Arg Leu Thr Phe Pro Glu Gln Ser Val Pro Gly Ala
 115 120 125
 Leu Gln Phe Arg Leu Thr Ser Val Asp Leu Glu Asp Trp Met Asp Val
 130 135 140
 Ser Leu Val Pro Ala Phe Asn Val Leu Gly Gln Ala Gly Ser Ala Val
 145 150 155 160
 Lys Pro Lys Pro Gln Val Tyr Ser Thr Leu Leu Asn Ser Gly Cys Gln
 165 170 175
 Gly Gly Glu His Ala Ala Cys Phe Thr Glu Leu Arg Arg Asn Phe Val
 180 185 190
 Asn Ile Arg Pro Ala Lys Leu Lys Asn Leu Ile Leu Leu Val Lys His
 195 200 205
 Trp Tyr His Gln Val Cys Leu Gln Gly Leu Trp Lys Glu Thr Leu Pro
 210 215 220
 Pro Val Tyr Ala Leu Glu Leu Leu Thr Ile Phe Ala Trp Glu Gln Gly
 225 230 235 240

Cys Lys Lys Asp Ala Phe Ser Leu Gly Glu Gly Leu Arg Thr Val Leu
 245 250 255

Gly Leu Ile Gln Gln His Gln His Leu Cys Val Phe Trp Thr Val Asn
 260 265 270

Tyr Gly Phe Glu Asp Pro Ala Val Gly Gln Phe Leu Gln Arg His Val
 275 280 285

Lys Arg Pro Arg Pro Val Ile Leu Asp Pro Ala Asp Pro Thr Trp Asp
 290 295 300

Leu Gly Asn Gly Ala Ala Trp His Trp Asp Leu His Ala Gln Glu Ala
 305 310 315 320

Ala Ser Cys Tyr Asp His Pro Cys Phe Leu Arg Gly Met Gly Asp Pro
 325 330 335

Val Gln Ser Trp Lys Gly Pro Gly Leu Pro Arg Ala Gly Cys Ser Gly
 340 345 350

Leu Gly His Pro Ile Gln Leu Asp Pro Asn Gln Lys Thr Pro Glu Asn
 355 360 365

Ser Lys Ser Leu Asn Ala Val Tyr Pro Arg Ala Gly Ser Lys Pro Pro
 370 375 380

Ser Cys Pro Ala Pro Gly Pro Thr Ala Glu Pro Ala Ser Tyr Pro Ser
 385 390 395 400

Val Pro Gly Met Ala Leu Asp Leu Ser Gln Ile Pro Thr Lys Glu Leu
 405 410 415

Asp Arg Phe Ile Gln Asp His Leu Lys Pro Ser Pro Gln Phe Gln Glu
 420 425 430

Gln Val Lys Lys Ala Ile Asp Ile Ile Leu Arg Cys Leu His Glu Asn
 435 440 445

Cys Val His Lys Ala Ser Arg Val Ser Lys Gly Gly Ser Phe Gly Arg
 450 455 460

Gly Thr Asp Leu Arg Asp Gly Cys Asp Val Glu Leu Ile Ile Phe Leu
 465 470 475 480

Asn Cys Phe Thr Asp Tyr Lys Asp Gln Gly Pro Arg Arg Ala Glu Ile
 485 490 495

Leu Asp Glu Met Arg Ala His Val Glu Ser Trp Trp Gln Asp Gln Val
 500 505 510

Pro Ser Leu Ser Leu Gln Phe Pro Glu Gln Asn Val Pro Glu Ala Leu
 515 520 525

Gln Phe Gln Leu Val Ser Thr Ala Leu Lys Ser Trp Thr Asp Val Ser
 530 535 540

Leu Leu Pro Ala Phe Asp Ala Val Gly Gln Leu Ser Ser Gly Thr Lys
 545 550 555 560

Pro Asn Pro Gln Val Tyr Ser Arg Leu Leu Thr Ser Gly Cys Gln Glu
 565 570 575

Gly Glu His Lys Ala Cys Phe Ala Glu Leu Arg Arg Asn Phe Met Asn
 580 585 590

Ile Arg Pro Val Lys Leu Lys Asn Leu Ile Leu Leu Val Lys His Trp
 595 600 605

Tyr Arg Gln Val Ala Ala Gln Asn Lys Gly Lys Gly Pro Ala Pro Ala
 610 615 620

Ser Leu Pro Pro Ala Tyr Ala Leu Glu Leu Leu Thr Ile Phe Ala Trp
 625 630 635 640

Glu Gln Gly Cys Arg Gln Asp Cys Phe Asn Met Ala Gln Gly Phe Arg
 645 650 655

Thr Val Leu Gly Leu Val Gln Gln His Gln Gln Leu Cys Val Tyr Trp
 660 665 670

Thr Val Asn Tyr Ser Thr Glu Asp Pro Ala Met Arg Met His Leu Leu
 675 680 685

Gly Gln Leu Arg Lys Pro Arg Pro Leu Val Leu Asp Pro Ala Asp Pro
 690 695 700

Thr Trp Asn Val Gly His Gly Ser Trp Glu Leu Leu Ala Gln Glu Ala

705		710		715		720
Ala Ala Leu Gly	Met Gln Ala Cys Phe Leu Ser Arg Asp Gly Thr Ser					
	725		730			735
Val Gln Pro Trp	Asp Val Met Pro Ala Leu Leu Tyr Gln Thr Pro Ala					
	740		745			750
Gly Asp Leu Asp	Lys Phe Ile Ser Glu Phe Leu Gln Pro Asn Arg Gln					
	755		760			765
Phe Leu Ala Gln	Val Asn Lys Ala Val Asp Thr Ile Cys Ser Phe Leu					
	770		775			780
Lys Glu Asn Cys	Phe Arg Asn Ser Pro Ile Lys Val Ile Lys Val Val					
	785		790		795	800
Lys Gly Gly Ser	Ser Ala Lys Gly Thr Ala Leu Arg Gly Arg Ser Asp					
	805		810			815
Ala Asp Leu Val	Val Phe Leu Ser Cys Phe Ser Gln Phe Thr Glu Gln					
	820		825			830
Gly Asn Lys Arg	Ala Glu Ile Ile Ser Glu Ile Arg Ala Gln Leu Glu					
	835		840			845
Ala Cys Gln Gln	Glu Arg Gln Phe Glu Val Lys Phe Glu Val Ser Lys					
	850		855			860
Trp Glu Asn Pro	Arg Val Leu Ser Phe Ser Leu Thr Ser Gln Thr Met					
	865		870		875	880
Leu Asp Gln Ser	Val Asp Phe Asp Val Leu Pro Ala Phe Asp Ala Leu					
	885		890			895
Gly Gln Leu Val	Ser Gly Ser Arg Pro Ser Ser Gln Val Tyr Val Asp					
	900		905			910
Leu Ile His Ser	Tyr Ser Asn Ala Gly Glu Tyr Ser Thr Cys Phe Thr					
	915		920			925
Glu Leu Gln Arg	Asp Phe Ile Ile Ser Arg Pro Thr Lys Leu Lys Ser					
	930		935			940

34

Leu Ile Arg Leu Val Lys His Trp Tyr Gln Gln Cys Thr Lys Ile Ser
 945 950 955 960

Lys Gly Arg Gly Ser Leu Pro Pro Gln His Gly Leu Glu Leu Leu Thr
 965 970 975

Val Tyr Ala Trp Glu Gln Gly Gly Lys Asp Ser Gln Phe Asn Met Ala
 980 985 990

Glu Gly Phe Arg Thr Val Leu Glu Leu Val Thr Gln Tyr Arg Gln Leu
 995 1000 1005

Cys Ile Tyr Trp Thr Ile Asn Tyr Asn Ala Lys Asp Lys Thr Val
 1010 1015 1020

Gly Asp Phe Leu Lys Gln Gln Leu Gln Lys Pro Arg Pro Ile Ile
 1025 1030 1035

Leu Asp Pro Ala Asp Pro Thr Gly Asn Leu Gly His Asn Ala Arg
 1040 1045 1050

Trp Asp Leu Leu Ala Lys Glu Ala Ala Ala Cys Thr Ser Ala Leu
 1055 1060 1065

Cys Cys Met Gly Arg Asn Gly Ile Pro Ile Gln Pro Trp Pro Val
 1070 1075 1080

Lys Ala Ala Val
 1085

<210> 11
 <211> 1104
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(1104)

<300>
 <301> Coccia, E.M., et al.
 <302> A full-length murine 2-5A synthetase cDNA transfected in NIH-3T3 cells
 <303> Virology
 <304> 179
 <305> 1
 <306> 228-233
 <307> 1990

35

<308> NCBI/M33863

<309> 1993-06-11

<400> 11

atg gag cac gga ctc agg agc atc cca gcc tgg acg ctg gac aag ttc	48
Met Glu His Gly Leu Arg Ser Ile Pro Ala Trp Thr Leu Asp Lys Phe	
1 5 10 15	
ata gag gat tac ctc ctt ccc gac acc acc ttt ggt gct gat gtc aaa	96
Ile Glu Asp Tyr Leu Leu Pro Asp Thr Thr Phe Gly Ala Asp Val Lys	
20 25 30	
tca gcc gtc aat gtc gtg tgt gat ttc ctg aag gag aga tgc ttc caa	144
Ser Ala Val Asn Val Val Cys Asp Phe Leu Lys Glu Arg Cys Phe Gln	
35 40 45	
ggt gct gcc cac cca gtg agg gtc tcc aag gtg gtg aag ggt ggc tcc	192
Gly Ala Ala His Pro Val Arg Val Ser Lys Val Val Lys Gly Gly Ser	
50 55 60	
tca ggc aaa ggc acc aca ctc aag ggc agg tca gac gct gac ctg gtg	240
Ser Gly Lys Gly Thr Thr Leu Lys Gly Arg Ser Asp Ala Asp Leu Val	
65 70 75 80	
gtg ttc ctt aac aat ctc acc agc ttt gag gat cag tta aac cga cgg	288
Val Phe Leu Asn Asn Leu Thr Ser Phe Glu Asp Gln Leu Asn Arg Arg	
85 90 95	
gga gag ttc atc aag gaa att aag aaa cag ctg tac gag gtt cag cat	336
Gly Glu Phe Ile Lys Glu Ile Lys Lys Gln Leu Tyr Glu Val Gln His	
100 105 110	
gag aga cgt ttt aga gtc aag ttt gag gtc cag agt tca tgg tgg ccc	384
Glu Arg Arg Phe Arg Val Lys Phe Glu Val Gln Ser Ser Trp Trp Pro	
115 120 125	
aac gcc cgg tct ctg agc ttc aag ctg agc gcc ccc cat ctg cat cag	432
Asn Ala Arg Ser Leu Ser Phe Lys Leu Ser Ala Pro His Leu His Gln	
130 135 140	
gag gtg gag ttt gat gtg ctg cca gcc ttt gat gtc ctg ggt cat gtt	480
Glu Val Glu Phe Asp Val Leu Pro Ala Phe Asp Val Leu Gly His Val	
145 150 155 160	
aat act tcc agc aag cct gat ccc aga atc tat gcc atc ctc atc gag	528
Asn Thr Ser Ser Lys Pro Asp Pro Arg Ile Tyr Ala Ile Leu Ile Glu	
165 170 175	
gaa tgt acc tcc ctg ggg aag gat ggc gag ttc tct acc tgc ttc acg	576
Glu Cys Thr Ser Leu Gly Lys Asp Gly Glu Phe Ser Thr Cys Phe Thr	
180 185 190	
gag ctc cag cgg aac ttc ctg aag cag cgc cca acc aag ctg aag agt	624
Glu Leu Gln Arg Asn Phe Leu Lys Gln Arg Pro Thr Lys Leu Lys Ser	
195 200 205	
ctc atc cgc ctg gtc aag cac tgg tac caa ctg tgt aag gag aag ctg	672
Leu Ile Arg Leu Val Lys His Trp Tyr Gln Leu Cys Lys Glu Lys Leu	
210 215 220	

36

ggg aag cca ttg cct cca cag tac gcc cta gag ttg ctc act gtc ttt 720
 Gly Lys Pro Leu Pro Pro Gln Tyr Ala Leu Glu Leu Leu Thr Val Phe
 225 230 235 240

gcc tgg gaa caa ggg aat gga tgt tat gag ttc aac aca gcc cag ggc 768
 Ala Trp Glu Gln Gly Asn Gly Cys Tyr Glu Phe Asn Thr Ala Gln Gly
 245 250 255

ttc cgg acc gtc ttg gaa ctg gtc atc aat tat cag cat ctt cga atc 816
 Phe Arg Thr Val Leu Glu Leu Val Ile Asn Tyr Gln His Leu Arg Ile
 260 265 270

tac tgg aca aag tat tat gac ttt caa cac cag gag gtc tcc aaa tac 864
 Tyr Trp Thr Lys Tyr Tyr Asp Phe Gln His Gln Glu Val Ser Lys Tyr
 275 280 285

ctg cac aga cag ctc aga aaa gcc agg cct gtg atc ctg gac cca gct 912
 Leu His Arg Gln Leu Arg Lys Ala Arg Pro Val Ile Leu Asp Pro Ala
 290 295 300

gac cca aca ggg aat gtg gcc ggt ggg aac cca gag ggc tgg agg cgg 960
 Asp Pro Thr Gly Asn Val Ala Gly Gly Asn Pro Glu Gly Trp Arg Arg
 305 310 315 320

ttg gct gaa gag gct gat gtg tgg cta tgg tac cca tgt ttt att aaa 1008
 Leu Ala Glu Glu Ala Asp Val Trp Leu Trp Tyr Pro Cys Phe Ile Lys
 325 330 335

aag gat ggt tcc cga gtg agc tcc tgg gat gtg ccg acg gtg gtt cct 1056
 Lys Asp Gly Ser Arg Val Ser Ser Trp Asp Val Pro Thr Val Val Pro
 340 345 350

gta cct ttt gag cag gta gaa gag aac tgg aca tgt atc ctg ctg tga 1104
 Val Pro Phe Glu Gln Val Glu Glu Asn Trp Thr Cys Ile Leu Leu
 355 360 365

<210> 12
 <211> 367
 <212> PRT
 <213> Mus musculus

<400> 12

Met Glu His Gly Leu Arg Ser Ile Pro Ala Trp Thr Leu Asp Lys Phe
 1 5 10 15

Ile Glu Asp Tyr Leu Leu Pro Asp Thr Thr Phe Gly Ala Asp Val Lys
 20 25 30

Ser Ala Val Asn Val Val Cys Asp Phe Leu Lys Glu Arg Cys Phe Gln
 35 40 45

Gly Ala Ala His Pro Val Arg Val Ser Lys Val Val Lys Gly Gly Ser
 50 55 60

Ser Gly Lys Gly Thr Thr Leu Lys Gly Arg Ser Asp Ala Asp Leu Val
65 70 75 80

Val Phe Leu Asn Asn Leu Thr Ser Phe Glu Asp Gln Leu Asn Arg Arg
85 90 95

Gly Glu Phe Ile Lys Glu Ile Lys Lys Gln Leu Tyr Glu Val Gln His
100 105 110

Glu Arg Arg Phe Arg Val Lys Phe Glu Val Gln Ser Ser Trp Trp Pro
115 120 125

Asn Ala Arg Ser Leu Ser Phe Lys Leu Ser Ala Pro His Leu His Gln
130 135 140

Glu Val Glu Phe Asp Val Leu Pro Ala Phe Asp Val Leu Gly His Val
145 150 155 160

Asn Thr Ser Ser Lys Pro Asp Pro Arg Ile Tyr Ala Ile Leu Ile Glu
165 170 175

Glu Cys Thr Ser Leu Gly Lys Asp Gly Glu Phe Ser Thr Cys Phe Thr
180 185 190

Glu Leu Gln Arg Asn Phe Leu Lys Gln Arg Pro Thr Lys Leu Lys Ser
195 200 205

Leu Ile Arg Leu Val Lys His Trp Tyr Gln Leu Cys Lys Glu Lys Leu
210 215 220

Gly Lys Pro Leu Pro Pro Gln Tyr Ala Leu Glu Leu Leu Thr Val Phe
225 230 235 240

Ala Trp Glu Gln Gly Asn Gly Cys Tyr Glu Phe Asn Thr Ala Gln Gly
245 250 255

Phe Arg Thr Val Leu Glu Leu Val Ile Asn Tyr Gln His Leu Arg Ile
260 265 270

Tyr Trp Thr Lys Tyr Tyr Asp Phe Gln His Gln Glu Val Ser Lys Tyr
275 280 285

Leu His Arg Gln Leu Arg Lys Ala Arg Pro Val Ile Leu Asp Pro Ala

38

290	295	300
Asp Pro Thr Gly Asn Val Ala Gly Gly Asn Pro Glu Gly Trp Arg Arg		
305	310	315 320
Leu Ala Glu Glu Ala Asp Val Trp Leu Trp Tyr Pro Cys Phe Ile Lys		
	325	330 335
Lys Asp Gly Ser Arg Val Ser Ser Trp Asp Val Pro Thr Val Val Pro		
	340	345 350
Val Pro Phe Glu Gln Val Glu Glu Asn Trp Thr Cys Ile Leu Leu		
355	360	365

<210> 13
 <211> 1590
 <212> DNA
 <213> Homo sapiens

<300>
 <301> Aissouni, Y. et al.
 <302> The cleavage/polyadenylation activity triggered by a U-rich motif
 sequence
 <303> J. Biol. Chem.
 <304> 277
 <305> 39
 <306> 35808-35814
 <307> 2002
 <308> NCBI/NM_016816
 <309> 2003-04-06

<300>
 <301> Behera, A.K., et al.
 <302> 2'-5' Oligoadenylate synthetase plays a critical role in
 interferon-gamma inhibition
 <303> J. Biol. Chem.
 <304> 277
 <305> 28
 <306> 25601-25608
 <307> 2002
 <308> NCBI/NM_016816
 <309> 2003-04-06

<300>
 <301> Sarkar, S.N., et al.
 <302> Identification of the substrate-binding sites of
 2'-5'-oligoadenylate synthetase
 <303> J. Biol. Chem.
 <304> 277
 <305> 27
 <306> 24321-24330
 <307> 2002
 <308> NCBI/NM_016816
 <309> 2003-04-06

- <300>
<301> Hovnanian, A., et al.
<302> The human 2',5'-oligoadenylate synthetase locus is composed of three distinct genes
<303> Genomics
<304> 52
<305> 3
<306> 267-277
<307> 1998
<308> NCBI/NM_016816
<309> 2003-04-06
- <300>
<301> Renault, B.
<302> A sequence-ready physical map of a region of 12q24.1
<303> Genomics
<304> 45
<305> 2
<306> 271-278
<307> 1997
<308> NCBI/NM_016816
<309> 2003-04-06
- <300>
<301> Nechiporuk, T., et al.
<302> A high-resolution PAC and BAC map of the SCA2 region
<303> Genomics
<304> 44
<305> 3
<306> 321-329
<307> 1997
<308> NCBI/NM_016816
<309> 2003-04-06
- <300>
<301> Wathelet, M.G., et al.
<302> Cloning and chromosomal location of human genes inducible by type I interferon
<303> Somat. Cell Mol. Genet.
<304> 14
<305> 5
<306> 415-426
<307> 1988
<308> NCBI/NM_016816
<309> 2003-04-06
- <300>
<301> Rutherford, M.N., et al.
<302> Interferon-induced binding of nuclear factors to promoter elements of the 2-5A synthetase gene
<303> EMBO J.
<304> 7
<305> 3
<306> 751-759
<307> 1988
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Wathelet, M.G., et al.
<302> New inducers revealed by the promoter sequence analysis of two
interferon-activated human genes
<303> Eur. J. Biochem.
<304> 169
<305> 2
<306> 313-321
<307> 1987
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Benech, P., et al.
<302> Interferon-responsive regulatory elements in the promoter of the
human 2',5'-oligo(A) synthetase gene
<303> Mol. Cell. Biol.
<304> 7
<305> 12
<306> 4498-4504
<307> 1987
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Hovanessian, A.G., et al.
<302> Identification of 69-kd and 100-kd forms of 2-5A synthetase
<303> EMBO J.
<304> 6
<305> 5
<306> 1273-1280
<307> 1987
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Williams, B.R., et al.
<302> Interferon-regulated human 2-5A synthetase gene maps to
chromosome
<303> Somat. Cell Mol. Genet.
<304> 12
<305> 4
<306> 403-408
<307> 1986
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Shiojiri, S., et al.
<302> Structure and expression of a cloned cDNA
<303> J. Biochem.
<304> 99
<305> 5
<306> 1455-1464
<307> 1986
<308> NCBI/NM_016816
<309> 2003-04-06

<300>

<301> Wathelet, M., et al.
<302> Full-length sequence and expression of the 42 kDa 2-5A synthetase
<303> FEBS Lett.
<304> 196
<305> 1
<306> 113-120
<307> 1986
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Benech, P., et al.
<302> Structure of two forms of the interferon-induced (2'-5') oligo A synthetase
<303> EMBO J.
<304> 4
<305> 9
<306> 2249-2256
<307> 1985
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Saunders, M.E., et al.
<302> Human 2-5A synthetase: characterization of a novel cDNA and corresponding gene structure
<303> EMBO J.
<304> 4
<305> 7
<306> 1761-1768
<307> 1985
<308> NCBI/NM_016816
<309> 2003-04-06

<300>
<301> Merlin, G., et al.
<302> Molecular cloning and sequence of partial cDNA for interferon-induced (2'-5') oligo(A) synthetase mRNA from human cells
<303> Proc. Natl. Acad. Sci. U.S.A.
<304> 80
<305> 16
<306> 4904-4908
<307> 1983
<308> NCBI/NM_016816
<309> 2003-04-06

<400> 13
gaggcagttc tgttgccact ctctctcctg tcaatgatgg atctcagaaa taccccagcc 60
aaatctctgg acaagttcat tgaagactat ctcttgccag acacgtgttt ccgcatgcaa 120
atcgaccatg ccattgacat catctgtggg ttcttgaagg aaaggtgctt ccgaggtagc 180
tctaccctg tgtgtgtgtc caaggtggta aaggggtggc cctcaggcaa gggcaccacc 240
ctcagaggcc gatctgacgc tgacctgggt gtcttctca gtctctcac cacttttcag 300
gatcagttaa atcgccgggg agagttcatc caggaaatta ggagacagct ggaagcctgt 360

```

caaagagaga gagcactttc cgtgaagttt gaggtccagg ctccacgctg gggcaacccc 420
cgtgcgctca gcttcgtact gagttcgctc cagctcgggg aggggggtgga gttcgatgtg 480
ctgcctgcct ttgatgccct gggtcagttg actggcagct ataaacctaa cccccaaatc 540
tatgtcaagc tcatcgagga gtgcaccgac ctgcagaaag agggcgagtt ctccacctgc 600
ttcacagaac tacagagaga cttcctgaag cagcgcccca ccaagctcaa gagcctcatc 660
cgcctagtca agcactggta ccaaaattgt aagaagaagc ttgggaagct gccacctcag 720
tatgccctgg agctcctgac ggtctatgct tgggagcgag ggagcatgaa aacacatttc 780
aacacagccc aaggatttcg gacggctcttg gaattagtca taaactacca gcaactctgc 840
atctactgga caaagtatta tgactttaaa aacccatta ttgaaaagta cctgagaagg 900
cagctcacga aaccagggcc tgtgatcctg gacccggcgg accctacagg aaacttgggt 960
ggtggagacc caaagggttg gaggcagctg gcacaagagg ctgaggcctg gctgaattac 1020
ccatgcttta agaattggga tgggtcccca gtgagctcct ggattctgct ggctgaaagc 1080
aacagtacag acgatgagac cgacgatccc aggacgtatc agaaatatgg ttacattgga 1140
acacatgagt accctcattt ctctcataga ccagcacgc tccaggcage atccacccca 1200
caggcagaag aggactggac ctgcaccatc ctctgaatgc cagtgcattc tgggggaaag 1260
ggctccagtg ttatctggac cagttccttc attttcaggt gggactcttg atccagagaa 1320
gacaaagctc ctcagtgagc tgggtgtataa tccaagacag aaccaagtc tcctgactcc 1380
tggccttcta tgccctctat cctatcatag ataacattct ccacagcctc acttcattcc 1440
acctattctc tgaaaatatt ccctgagaga gaacagagag atttagataa gagaatgaaa 1500
ttccagcctt gactttcttc tgtgcacctg atgggagggt aatgtctaata gtattatcaa 1560
taacaataaa aataaagcaa ataccaaaaa 1590

```

<210> 14

<211> 3068

<212> DNA

<213> Homo sapiens

<300>

<301> Hovnanian, A., et al.

<302> The human 2',5'-oligoadenylate synthetase locus is composed of three distinct genes

<303> Genomics

<304> 52

<305> 3

<306> 267-277

<307> 1998

<308> NCBI/NM_002535

<309> 2003-04-03

<300>

<301> Marie, I. and Hovanessian, A.G.

<302> The 69-kDa 2-5A synthetase is composed of two homologous and adjacent functional domains

<303> J. Biol. Chem.

<304> 267

<305> 14

<306> 9933-9939

<307> 1992

<308> NCBI/NM_002535

<309> 2003-04-03

<300>

<301> Marie, I., et al.

<302> Differential expression and distinct structure of 69- and 100-kDa forms of 2-5A synthetase

<303> J. Biol. Chem.

<304> 265

<305> 30

<306> 18601-18607

<307> 1990

<308> NCBI/NM_002535

<309> 2003-04-03

<300>

<301> Marie, I., et al.

<302> Preparation and characterization of polyclonal antibodies

<303> Biochem. Biophys. Res. Commun.

<304> 160

<305> 2

<306> 580-587

<307> 1989

<308> NCBI/NM_002535

<309> 2003-04-03

<300>

<301> Hovanessian, A.G., et al.

<302> Characterization of 69- and 100-kDa forms of 2-5A-synthetase

<303> J. Biol. Chem.

<304> 263

<305> 10

<306> 4959

<307> 1988

<308> NCBI/NM_002535

<309> 2003-04-03

<300>

<301> Hovanessian, A.G., et al.

<302> Identification of 69-kd and 100-kd forms of 2-5A synthetase

<303> EMBO J.

<304> 6

<305> 5

<306> 1273-1280

<307> 1987

<308> NCBI/NM_002535

<309> 2003-04-03

<400> 14
 cggcagccag ctgagagcaa tgggaaatgg ggagtcccag ctgtcctcgg tgcctgctca 60
 gaagctgggt tggtttatcc aggaatacct gaagccctac gaagaatgtc agacactgat 120
 cgacgagatg gtgaacacca tctgtgacgt ctgcaggaac cccgaacagt tccccctggg 180
 gcaggggagtg gccatagggtg gctcctatgg acggaaaaca gtcttaagag gcaactccga 240
 tggtagcctt gtccttttct tcagtgaactt aaaacaattc caggatcaga agagaagcca 300
 acgtgacatc ctcgataaaa ctggggataa gctgaagttc tgtctgttca cgaagtgggt 360
 gaaaaacaat ttcgagatcc agaagtcctt tgatgggtcc accatccagg tgttcacaaa 420
 aaatcagaga atctctttcg aggtgctggc cgccttcaac gctctgagct taaatgataa 480
 tcccagcccc tggatctatc gagagctcaa aagatccttg gataagacaa atgccagtcc 540
 tggtagtgtt gcagtctgct tcaactgaact ccagcagaag ttttttgaca accgtcctgg 600
 aaaactaaag gatattgatcc tcttgataaa gcaactggcat caacagtgcc agaaaaaat 660
 caaggattta ccctcgctgt ctccgtatgc cctggagctg cttacggtgt atgcctggga 720
 acaggggtgc agaaaagaca actttgacat tgctgaaggc gtcagaacgg ttctggagct 780
 gatcaaatgc caggagaagc tgtgtatcta ttggatggtc aactacaact ttgaagatga 840
 gaccatcagg aacatcctgc tgcaccagct ccaatcagcg aggccagtaa tcttggatcc 900
 agttgaccca accaataatg tgagtggaga taaaatatgc tggcaatggc tgaaaaaaga 960
 agctcaaacc tggttgactt ctcccaacct ggataatgag ttacctgcac catcttgga 1020
 tgtcctgcct gcaccactct tcacgacccc aggccacctt ctggataagt tcatcaagga 1080
 gtttctccag cccaacaaat gcttcctaga gcagattgac agtgctgtta acatcatccg 1140
 tacattcctt aaagaaaact gcttcgaca atcaacagcc aagatccaga ttgtccgggg 1200
 aggatcaacc gccaaaggca cagctctgaa gactggctct gatgccgatc tcgtcgtgtt 1260
 ccataactca cttaaaagct acacctcca aaaaaacgag cggcacaaaa tcgtcaagga 1320
 aatccatgaa cagctgaaag ccttttgagg ggagaaggag gaggagcttg aagtcagctt 1380
 tgagcctccc aagtggaagg ctcccagggt gctgagcttc tctctgaaat ccaaagtcct 1440
 caacgaaagt gtcagctttg atgtgcttcc tgcctttaat gcaactgggtc agctgagttc 1500
 tggctccaca ccagccccg aggtttatgc agggctcatt gatctgtata aatcctcgga 1560
 cctcccggga ggagagtttt ctacctgttt cacagtcctg cagcgaaact tcattcgctc 1620
 ccggcccacc aaactaaagg atttaattcg cctgggtgaag cactggtaca aagagtgtga 1680
 aaggaaactg aagccaaagg ggtctttgcc ccaaagtat gccttgagc tgctcaccat 1740

45

```

ctatgcctgg gagcagggga gtggagtgcc ggattttgac actgcagaag gtttccggac 1800
agtcctggag ctggtcacac aatatcagca gctcggcatc ttctggaagg tcaattacaa 1860
ctttgaagat gagaccgtga ggaagtttct actgagccag ttgcagaaaa ccaggcctgt 1920
gatcttggac ccaggcgaac ccacaggtga cgtgggtgga ggggaccgtt ggtgttggca 1980
tcttctggac aaagaagcaa aggttaggtt atcctctccc tgcttcaagg atgggactgg 2040
aaaccaata ccaccttga aagtgccggt aaaagtcac taaaggaggc gttgtctgga 2100
aatagccctg taacaggctt gaatcaaaga acttctccta ctgtagcaac ctgaaattaa 2160
ctcagacaca aataaaggaa acccagctca caggagctta aacagctggt cagccccct 2220
aagccccac tacaagtgat cctcaggcag gtaacccag attcatgcac tgtagggctg 2280
ggcgagcat ccctaggtct ctaccagta gatgccacta gccctcctct ccagtgaca 2340
acaaaagtc ttcacatgtt caaacgttcc cctgggttca cagatcttct tgcctttggc 2400
ttttggctcc accctcttta gctgttaatt tgagtactta tggccctgaa agcggccacg 2460
gtgcctccag atggcaggtt tgcaatccaa gcaggaagaa ggaaaagata cccaaaggtc 2520
aagaacacag tgattttatt agaagtttca tccgcaaatt ttcttccatt tcattgctca 2580
gaatgtcatg tggttacctg taacttgaag gtggctacaa agatgactgt ggaggtggtt 2640
gcacttgcca cccaaggatg tctgccacac ctctccaagc cctcctacct accaagatat 2700
acctgatata tccaccagat atctctcag atatacttgg ttctctccac caggttcttt 2760
ctttaaagca ggattctcaa ctttgatact tactcacatt gggctagaca gttctttggt 2820
tggaggtctc cttgtgcatg taggatgttg agcagcatgt gtggcctgta ccagtacat 2880
gccaccagt tgtgacaatt aaaagtgtct tgagacttta tcatgtgtct tctgccctag 2940
gtgagaacct ttgcactaca ggaacctac acccaacctg gggggaatgt agggaagagg 3000
tgccaagcca accgtggggt tagctctaata tattaagtta tgcattataa ataaatacca 3060
aaaaattg 3068

```

```

<210> 15
<211> 6270
<212> DNA
<213> Homo sapiens

```

```

<300>
<301> Rebouillat, D., et al.
<302> The 100-kDa 2',5'-oligoadenylate synthetase catalyzing
preferentially the synthesis of dimeric pppA2'p5'A molecules
<303> J. Biol. Chem.
<304> 274
<305> 3

```


<306> 1557-1565
 <307> 1999
 <308> NCBI/AF063613
 <309> 1999-05-04

 <300>
 <301> Rebouillat, D., and Hovanessian, A.G.
 <302> Direct Submission
 <303> Submitted (07-May-1998) Dept. of AIDS and Retroviruses, Institut Pasteur
 <304> 0
 <305> 0
 <306> 0
 <307> 1998
 <308> NCBI/AF063613
 <309> 1999-05-04

 <400> 15
 gccctgcttc cccttgcacc tgcgcgcggc ggccatggac ttgtacagca ccccgccgc 60
 tgcgctggac aggttcgtgg ccagaaggct gcagccgcgg aaggagttcg tagagaaggc 120
 gcggcgcgct ctgggcgccc tggccgctgc cctgagggag cgcgggggccc gcctcggtgc 180
 tgctgccccg cgggtgctga aaactgtcaa gggaggctcc tcgggcgggg gcacagctct 240
 caagggtggc tgtgattctg aacttgtcat cttcctcgac tgcttcaaga gctatgtgga 300
 ccagagggcc cgccgtgcag agatcctcag tgagatgcgg gcatcgctgg aatcctggtg 360
 gcagaaccca gtccctggtc tgagactcac gtttcctgag cagagcgtgc ctggggccct 420
 gcagttccgc ctgacatccg tagatcttga ggactggatg gatgttagcc tgggtgcctgc 480
 cttcaatgtc ctgggtcagg ccggctccgc ggtcaaacc aagccacaag tctactctac 540
 cctcctcaac agtggctgcc aagggggcga gcatgcggcc tgcttcacag agctgcggag 600
 gaactttgtg aacattcgcc cagccaagtt gaagaacct atcttgctgg tgaagcactg 660
 gtaccaccag gtgtgcctac aggggttgtg gaaggagacg ctgcccccg tctatgccct 720
 ggaattgctg accatcttcg cctgggagca gggctgtaag aaggatgctt tcagcctagg 780
 cgaaggcctc cgaactgtcc tgggcctgat ccaacagcat cagcacctgt gtgttttctg 840
 gactgtcaac tatggcttcg aggaccctgc agttgggcag ttcttgacgc ggcacgttaa 900
 gagaccagc cctgtgatcc tggaccagc tgaccccaca tgggacctgg ggaatggggc 960
 agcctggcac tgggatttgc atgcccagga ggcagcatcc tgctatgacc acccatgctt 1020
 tctgaggggg atgggggacc cagtgcagtc ttggaagggg ccgggccttc cacgtgctgg 1080
 atgctcaggt ttgggccacc ccatccagct agaccctaac cagaagaccc ctgaaaacag 1140
 caagagcctc aatgctgtgt acccaagagc agggagcaaa cctccctcat gccagctcc 1200
 tggccccact gcggagccag catcgtaccc ctctgtgccg ggaatggcct tggacctgtc 1260

tcagatcccc	accaaggagc	tggaccgctt	catccaggac	cacctgaagc	cgagccccc	1320
gttccaggag	caggtgaaaa	aggccatcga	catcatcttg	cgctgcctcc	atgagaactg	1380
tgttcacaag	gcctcaagag	tcagtaaagg	gggctcattt	ggccggggca	cagacctaa	1440
ggatggctgt	gatgttgaac	tcacatctt	cctcaactgc	ttcacggact	acaaggacca	1500
ggggccccc	cgcgagaga	tccttgatga	gatgagagc	cacgtagaat	cctgggtggc	1560
ggaccagggt	cccagcctga	gccttcagtt	tcctgagcag	aatgtgcctg	aggctctgca	1620
gttccagctg	gtgtccacag	ccctgaagag	ctggacggat	gttagcctgc	tgctgcctt	1680
cgatgctgtg	gggcagctca	gttctggcac	caaaccaaat	ccccaggtct	actcgaggct	1740
cctcaccagt	ggctgccagg	agggcgagca	taaggcctgc	ttcgagagc	tgcgaggagaa	1800
cttcatgaac	attcgccctg	tcaagctgaa	gaacctgatt	ctgctggtga	agcactggta	1860
ccgccagggt	gcggctcaga	acaaaggaaa	aggaccagcc	cctgcctctc	tgccccagc	1920
ctatgccctg	gagctcctca	ccatctttgc	ctgggagcag	ggctgcaggc	aggattgttt	1980
caacatggcc	caaggcttcc	ggacggtgct	ggggctcgtg	caacagcatc	agcagctctg	2040
tgtctactgg	acggtcaact	atagcactga	ggaccagcc	atgagaatgc	accttcttgg	2100
ccagcttcga	aaaccagac	ccctggctct	ggaccccgct	gatccacct	ggaacgtggg	2160
ccacggtagc	tgggagctgt	tggcccagga	agcagcagcg	ctggggatgc	aggcctgctt	2220
tctgagtaga	gacgggacat	ctgtgcagcc	ctgggatgtg	atgccagccc	tcctttacca	2280
aacccagct	ggggaccttg	acaagttcat	cagtgaattt	ctccagccca	accgccagtt	2340
cctggcccag	gtgaacaagg	ccgttgatac	catctgttca	tttttgaagg	aaaactgctt	2400
ccggaattct	cccatcaaag	tgatcaagggt	ggtaagggt	ggctcttcag	ccaaaggcac	2460
agctctgcga	ggccgctcag	atgccgacct	cgtggtgttc	ctcagctgct	tcagccagtt	2520
cactgagcag	ggcaacaagc	gggcccagat	catctccgag	atccgagccc	agctggaggc	2580
atgtcaacag	gagcggcagt	tcgaggtcaa	gtttgaagtc	tccaaatggg	agaatccccg	2640
cgtgctgagc	ttctcactga	catcccagac	gatgctggac	cagagtgtgg	actttgatgt	2700
gctgccagcc	tttgacgccc	taggccagct	ggtctctggc	tccaggccca	gctctcaagt	2760
ctacgtcgac	ctcatccaca	gctacagcaa	tgccggcgag	tactccacct	gcttcacaga	2820
gctacaacgg	gacttcatca	tctctcgccc	taccaagctg	aagagcctga	tccggctgggt	2880
gaagcactgg	taccagcagt	gtaccaagat	ctccaagggg	agaggctccc	tacccccaca	2940
gcacgggctg	gaactcctga	ctgtgtatgc	ctgggagcag	ggcgggaagg	actcccagtt	3000

caacatggct	gagggcttcc	gcacggtcct	ggagctggtc	accaggtacc	gccagctctg	3060
tatctactgg	accatcaact	acaacgcaa	ggacaagact	gttggagact	tctgaaaca	3120
gcagcttcag	aagcccaggc	ctatcatcct	ggatccggct	gacccgacag	gcaacctggg	3180
ccacaatgcc	cgctgggacc	tgctggccaa	ggaagctgca	gcctgcacat	ctgccctgtg	3240
ctgcatggga	cggaatggca	tcccatacca	gccatggcca	gtgaaggctg	ctgtgtgaag	3300
ttgagaaaat	cagcggctct	actggatgaa	gagaagatgg	acaccagccc	tcagcatgag	3360
gaaattcagg	gtcccctacc	agatgagaga	gattgtgtac	atgtgtgtgt	gagcacatgt	3420
gtgcatgtgt	gtgcacacgt	gtgcatgtgt	gtgttttagt	gaatctgctc	tcccagctca	3480
cacactcccc	tgccctccat	ggcttacaca	ctaggatcca	gactccatgg	tttgacacca	3540
gcctgcgttt	gcagcttctc	tgtaacttcc	atgactctat	cctcatacca	ccactgctgc	3600
ttcccacca	gctgagaatg	ccccctcctc	cctgactcct	ctctgcccac	gcaaattagc	3660
tcacatcttt	cctcctgctg	caatccatcc	cttcctccca	ttggcctctc	cttgccaaat	3720
ctaaatactt	tatatagggg	tggcagagag	ttcccatctc	atctgtcagc	cacagtcatt	3780
tggtactggc	tacctggagc	cttatcttct	gaagggtttt	aaagaatggc	caattagctg	3840
agaagaatta	tctaataaat	tagtgatgtc	tgccatggat	gcagtagagg	aaagtgggtg	3900
tacaagtgcc	atgattgatt	agcaatgtct	gcactggata	tggaaaaaag	aaggtgcttg	3960
cagggtttaca	gtgtatatgt	gggctattga	agagccctct	gagctcgggt	gctagcagga	4020
gagcatgccc	atattggctt	actttgtctg	ccacagacac	agacagaggg	agttgggaca	4080
tgcatgctat	ggggaccctc	ttgttggaca	cctaattgga	tgccctcttc	tgagaggcct	4140
ccttttcttc	accttttatg	ctgcactcct	cccctagttt	acacatcttg	atgctgtggc	4200
tcagtttgcc	ttcctgaatt	tttattgggt	ccctgttttc	tctcctaaca	tgctgagatt	4260
ctgcatcccc	acagcctaaa	ctgagccagt	ggccaaacaa	ccgtgctcag	cctgtttctc	4320
tctgccctct	agagcaaggc	ccaccaggtc	catccaggag	gctctcctga	cctcaagtcc	4380
aacaacagtg	tccacactag	tcaaggttca	gccagaaaa	cagaaagcac	tctaggaatc	4440
ttaggcagaa	agggatttta	tctaaatcac	tggaaaggct	ggaggagcag	aaggcagagg	4500
ccaccactgg	actattgggt	tcaatattag	accactgtag	ccgaatcaga	ggccagagag	4560
cagccactgc	tactgcta	gccaccacta	cccctgccat	cactgcccc	catggacaaa	4620
actggagtcg	agacctaggt	tagattcctg	caaccacaaa	catccatcag	ggatggccag	4680
ctgccagagc	tgcggaaga	cggatcccac	ctccctttct	tagcagaatc	taaattacag	4740
ccagacctct	ggctgcagag	gagtcctgaga	catgtatgat	tgaatgggtg	ccaagtgcc	4800

```

ggggggcggag tccccagcag atgcatcctg gccatctgtt gcgtggatga gggagtgggt 4860
ctatctcaga ggaaggaaca ggaaacaaag aaaggaagcc actgaacatc ccttctctgc 4920
tccacaggag tgtcttagac agcctgactc tccacaaacc actgttaaaa cttacctgct 4980
aggaatgcta gattgaatgg gatgggaaga gccttccctc attattgtca ttcttggaga 5040
gaggtgagca accaaggga gctcctctga ttcacctaga acctgttctc tgccgtcttt 5100
ggctcagcct acagagacta gagtaggtga agggacagag gacagggtt ctaataacctg 5160
tgccatattg acagcctcca tccctgtccc ccatcttggt gctgaaccaa cgctaagggc 5220
accttcttag actcacctca tcgatactgc ctggtaatcc aaagctagaa ctctcaggac 5280
cccaaactcc acctcttgga ttggccctgg ctgctgccac acacatatcc aagagctcag 5340
ggccagttct ggtgggcagc agagacctgc tctgccaaagt tgtccagcag cagagtggcc 5400
ctggcctggg catcacaagc cagtgatgct cctgggaaga ccaggtggca ggtcgcagtt 5460
gggtaccttc cattcccacc acacagactc tgggcctccc cgcaaatgg ctccagaatt 5520
agagtaatta tgagatggtg ggaaccagag caactcaggt gcatgataca aggagagggt 5580
gtcatctggg tagggcagag aggagggtt gctcatctga acaggggtgt atttcattcc 5640
aggccctcag tctttggcaa tggccaccct ggtgttgga tattggcccc actgtaactt 5700
ttgggggctt cccggtctag ccacaccctc ggatggaaag acttgactgc ataaagatgt 5760
cagttctccc tgagttgatt gataggctta atggtcaccc taaaaacacc cacatatgct 5820
tttcgatgga accagataag ttgacgctaa agttcttatg gaaaaataca cagcaatag 5880
ctaggaaaac acagggaaag aagagttctg agcagggcct agtcttagcc aatattaaaa 5940
catactatga agcctctgat acttaaacag catggcgctg gtacgtaaata agaccaatgc 6000
agttaggtgg ctctttccaa gactctgggg aaaaaagtag taaaaagcta aatgcaatca 6060
atcagcaatt gaaagctaag tgagagagcc agagggcctc cttggtggta aaagagggtt 6120
gcatttcttg cagccagaag gcagagaaag tgaagaccaa gtccagaact gaatcctaag 6180
aaatgcagga ctgcaaagaa attggtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtttaa 6240
tttttaaaaa gtttttattc ggaatccgcg 6270

```

```

<210> 16
<211> 1412
<212> DNA
<213> Mus musculus

```

```

<300>
<301> Coccia, E.M., et al.

```

<302> A full-length murine 2-5A synthetase cDNA transfected in NIH-3T3
 cells
 <303> Virology
 <304> 179
 <305> 1
 <306> 228-233
 <307> 1990
 <308> NCBI/M33863
 <309> 1993-06-11

 <400> 16
 ccaggctggg agaccagga agctccagac ttagcatgga gcacggactc aggagcatcc 60
 cagcctggac gctggacaag ttcatagagg attacctcct tcccgacacc acctttgggtg 120
 ctgatgtcaa atcagccgtc aatgtcgtgt gtgatttcct gaaggagaga tgcttccaag 180
 gtgctgcccc cccagtgagg gtctccaagg tggatgaagg tggctcctca ggcaaaggca 240
 ccacactcaa gggcagggtc gacgctgacc tgggtggtgtt ccttaacaat ctcaccagct 300
 ttgaggatca gttaaaccga cggggagagt tcatcaagga aattaagaaa cagctgtacg 360
 aggttcagca tgagagacgt tttagagtca agtttgagggt ccagagttca tgggtggcccc 420
 acgcccgggtc tctgagcttc aagctgagcg cccccatct gcatcaggag gtggagtttg 480
 atgtgctgcc agcctttgat gtcttgggtc atgttaatac ttccagcaag cctgatcccc 540
 gaatctatgc catcctcatc gaggaatgta cctccctggg gaaggatggc gagttctcta 600
 cctgcttcac ggagctccag cggaacttcc tgaagcagcg cccaaccaag ctgaagagtc 660
 tcatccgcct ggtcaagcac tggtagcaac tgtgtaagga gaagctgggg aagccattgc 720
 ctccacagta cgccctagag ttgctcactg tctttgcctg ggaacaaggg aatggatggt 780
 atgagttcaa cacagcccag ggcttccgga ccgtcttgga actgggtcatc aattatcagc 840
 atcttcgaat ctactggaca aagtattatg actttcaaca ccaggagggtc tccaaatacc 900
 tgcacagaca gctcagaaaa gccaggcctg tgatcctgga ccagctgac ccaacaggga 960
 atgtggccgg tgggaaccca gagggctgga ggcgggtggc tgaagaggct gatgtgtggc 1020
 tatggtaccc atgttttatt aaaaaggatg gttcccgagt gagctcctgg gatgtgccga 1080
 cgggtgggtcc tgtacctttt gagcaggtag aagagaactg gacatgtatc ctgctgtgag 1140
 cacagcagca cctgcccagg agactgctgg tcaggggcat ttgctgctct gctgcaggcc 1200
 catgaccagc tgagggaggg cccacactgg catcagactc cgtgcttctg atgctgcca 1260
 gccatgtttg actcctgtcc aatcacagcc agccttcctc aacagattca gaaggagagg 1320
 aaagaacaca cgcttggtgt ccatctgtcc acctgttgga aggttctgtc tgacaaagtc 1380
 tgatcaacaa taaaccacag cagggtgccgt ca 1412

<210> 17
 <211> 1193
 <212> DNA
 <213> Homo sapiens

<400> 17
 tgaagatcag ctattagaag agaaagatca gttaagtcct ttggacctga tcagcttgat 60
 acaagaacta ctgatttcaa cttcttttggc ttaattctct cggaaacgat gaaatataca 120
 agttatatct tggcttttca gctctgcacg gttttgggtt ctcttggctg ttactgccag 180
 gacccatatg taaaagaagc agaaaacctt aagaaatatt ttaatgcagg tcattcagat 240
 gtagcggata atggaactct tttcttaggc attttgaaga attggaaaga ggagagtgc 300
 agaaaaataa tgcagagcca aattgtctcc ttttacttca aactttttta aaacttttaa 360
 gatgaccaga gcatccaaaa gagtgtggag accatcaagg aagacatgaa tgtcaagttt 420
 ttcaatagca acaaaaagaa acgagatgac ttcgaaaagc tgactaatta ttcggtaact 480
 gacttgaatg tccaacgcaa agcaatacat gaactcatcc aagtgatggc tgaactgtcg 540
 ccagcagcta aaacagggaa gcgaaaaagg agtcagatgc tgtttcaagg tcgaagagca 600
 tcccagtaat gggtgtcctg cctgcaatat ttgaatttta aatctaaatc tatttattaa 660
 tatttaacat tatttatatg gggaaatatat ttttagactc atcaatcaaa taagtattta 720
 taatagcaac ttttgtgtaa tgaaaatgaa tatctattaa tatatgtatt atttataatt 780
 cctatatcct gtgactgtct cacttaatcc tttgttttct gactaattag gcaaggctat 840
 gtgattacaa ggcttttatct cagggggccaa ctaggcagcc aacctaagca agatcccatg 900
 gggtgtgtgt ttatttcact tgatgataca atgaacactt ataagtgaag tgatactatc 960
 cagttactgc cggtttgaaa atatgcctgc aatctgagcc agtgctttta tggcatgtca 1020
 gacagaactt gaatgtgtca ggtgaccctg atgaaaacat agcatctcag gagatttcat 1080
 gcctgggtgct tccaaatatt gttgacaact gtgactgtac ccaaattggaa agtaactcat 1140
 ttgttaaaat tatcaatatc taatatatat gaataaagtg taagttcaca act 1193

<210> 18
 <211> 166
 <212> PRT
 <213> Homo sapiens

<400> 18

Met Lys Tyr Thr Ser Tyr Ile Leu Ala Phe Gln Leu Cys Ile Val Leu
 1 5 10 15

52

Gly Ser Leu Gly Cys Tyr Cys Gln Asp Pro Tyr Val Lys Glu Ala Glu
 20 25 30

Asn Leu Lys Lys Tyr Phe Asn Ala Gly His Ser Asp Val Ala Asp Asn
 35 40 45

Gly Thr Leu Phe Leu Gly Ile Leu Lys Asn Trp Lys Glu Glu Ser Asp
 50 55 60

Arg Lys Ile Met Gln Ser Gln Ile Val Ser Phe Tyr Phe Lys Leu Phe
 65 70 75 80

Lys Asn Phe Lys Asp Asp Gln Ser Ile Gln Lys Ser Val Glu Thr Ile
 85 90 95

Lys Glu Asp Met Asn Val Lys Phe Phe Asn Ser Asn Lys Lys Lys Arg
 100 105 110

Asp Asp Phe Glu Lys Leu Thr Asn Tyr Ser Val Thr Asp Leu Asn Val
 115 120 125

Gln Arg Lys Ala Ile His Glu Leu Ile Gln Val Met Ala Glu Leu Ser
 130 135 140

Pro Ala Ala Lys Thr Gly Lys Arg Lys Arg Ser Gln Met Leu Phe Gln
 145 150 155 160

Gly Arg Arg Ala Ser Gln
 165

<210> 19

<211> 838

<212> DNA

<213> Homo sapiens

<400> 19

gagtctaact gcaacctttc gaagcctttg ctctggcaca acaggtagta ggcgacactg 60
 gtcgtgttgt tgacatgacc aacaagtgtc tcttccaaat tgctctcctg ttgtgcttct 120
 ccacgacagc tctttccatg agctacaact tgcttggatt cctacaaaga agcagcaatt 180
 gtcagtgtca gaagctcctg tggcaattga atgggaggct tgaatactgc ctcaaggaca 240
 ggaggaactt tgacatccct gaggagatta agcagctgca gcagttccag aaggaggacg 300
 ccgcagtgac catctatgag atgctccaga acatctttgc tattttcaga caagattcat 360
 cgagcactgg ctggaatgag actattgttg agaacctcct ggctaattgc tatcatcaga 420

gaaaccatct gaagacagtc ctggaagaaa aactggagaa agaagatttc accaggggaa 480
 aacgcatgag cagtctgcac ctgaaaagat attatgggag gattctgcat tacctgaagg 540
 ccaaggagga cagtcactgt gcctggacca tagtcagagt ggaaatccta aggaactttt 600
 acgtcattaa cagacttaca ggttacctcc gaaactgaag atctcctagc ctgtgcctct 660
 gggacgggac aattgcttca agcattcttc aaccagcaga tgctgtttta gtgactgatg 720
 gcgaatgtac tgcatatgaa aggacactag aagattttga aatttttatt aaattatgag 780
 gtatttttat ttattttaat tttatttttg aaaataaatt atttttggtg caaaagtc 838

<210> 20
 <211> 840
 <212> DNA
 <213> Homo sapiens

<400> 20
 acattctaac tgcaaccttt cgaagccttt gctctggcac aacaggtagt aggcgacact 60
 gttcgtgttg tcaacatgac caacaagtgt ctctccaaa ttgctctcct gttgtgcttc 120
 tccactacag ctctttccat gagctacaac ttgcttggat tcctacaaag aagcagcaat 180
 tttcagtgtc agaagctcct gtggcaattg aatgggaggc ttgaatactg cctcaaggac 240
 aggatgaact ttgacatccc tgaggagatt aagcagctgc agcagttcca gaaggaggac 300
 gccgcattga ccatctatga gatgctccag aacatctttg ctattttcag acaagattca 360
 tctagcactg gctggaatga gactattggt gagaacctcc tggctaattg ctatcatcag 420
 ataaaccatc tgaagacagt cctggaagaa aaactggaga aagaagattt caccagggga 480
 aaactcatga gcagtctgca cctgaaaaga tattatggga ggattctgca ttacctgaag 540
 gccaaaggagt acagtcactg tgcttgacc atagtcagag tggaaatcct aaggaacttt 600
 tacttcatta acagacttac aggttacctc cgaaactgaa gatctcctag cctgtgcctc 660
 tgggactgga caattgcttc aagcattctt caaccagcag atgctgttta agtgactgat 720
 ggctaattgta ctgcatatga aaggacacta gaagattttg aaatttttat taaattatga 780
 gttattttta tttattttaa ttttattttg gaaaataaat tatttttggg gcaaaagtca 840

<210> 21
 <211> 588
 <212> DNA
 <213> Homo sapiens

<400> 21
 atggccttcg tgctctctct actcatggcc ctgggtgctgg tcagctacgg cccaggagga 60

tccctgggtt gtgacctgtc tcagaaccac gtgctgggtg gcaggaagaa cctcaggctc 120
 ctggacgaaa tgaggagact ctcccctcac ttttgtctgc aggacagaaa agacttcgct 180
 ttaccccagg aaatggtgga gggcgggccag ctccaggagg cccaggccat ctctgtgctc 240
 catgagatgc tccagcagag cttcaacctc ttccacacag agcactcctc tgctgcctgg 300
 gacaccaccc tcctggagcc atgccgcact ggactccatc agcagctgga caacctggat 360
 gcctgcctgg ggcaggtgat gggagaggaa gactctgccc tgggaaggac gggccccacc 420
 ctggctctga agaggtactt ccagggcatc catgtctacc tgaaagagaa gggatacagc 480
 gactgcgctt gggaaaccgt cagactggaa atcatgagat ccttctcttc attaatacagc 540
 ttgcaagaaa ggttaagaat gatggatgga gacctgagct caccttga 588

<210> 22
 <211> 961
 <212> DNA
 <213> Homo sapiens

<400> 22
 tgagcctaaa ccttaggctc acccatittca accagtctag cagcatctgc aacatctaca 60
 atggccttga cctttgcttt actggtggcc ctccctggtgc tcagctgcaa gtcaagctgc 120
 tctgtgggct gtgatctgcc tcaaaccac agcctgggta gcaggaggac cttgatgctc 180
 ctggcacaga tgaggaaaat ctctcttttc tcctgcttga aggacagaca tgactttgga 240
 tttccccagg aggagtgttg caaccagttc caaaaggctg aaaccatccc tgtcctccat 300
 gagatgatcc agcagatctt caatctcttc agcacaaagg actcatctgc tgcttgggat 360
 gagaccctcc tagacaaatt ctacactgaa ctctaccagc agctgaatga cctggaagcc 420
 tgtgtgatac aggggggtgg ggtgacagag actcccctga tgaaggagga ctccattctg 480
 gctgtgagga aatacttcca aagaatcact ctctatctga aagagaagaa atacagccct 540
 tgtgcctggg aggttgtcag agcagaaatc atgagatctt tttctttgtc aacaaacttg 600
 caagaaagtt taagaagtaa ggaatgaaaa ctggttcaac atggaaatga ttttcattaa 660
 ttcgtatgcc agctcacctt tttatgatct gccatttcaa agactcatgt ttctgctatg 720
 accatgacac gatttaaata tttttcaaata gtttttagga gtattaatca acattgtatt 780
 cagctcttaa ggcactagtc cttacagag gaccatgctg actgatccat tatctattta 840
 aatattttta aaatattatt tatttaacta tttataaaaac aacttatttt tgttcatatt 900
 acgtcatgtg cacctttgca cagtgggtta tgtaataaaa tatgttcttt gtatttggtg 960
 a 961

<210> 23
 <211> 737
 <212> DNA
 <213> Homo sapiens

<400> 23
 taccacacctc aggtagccta gtgatatattg caaaatccca atggcccgggt ccttttcttt 60
 actgatggcc gtgctggtac tcagctacaa atccatctgc tctctgggct gtgatctgcc 120
 tcagacccac agcctgcgta ataggagggc cttgatactc ctggcacaaa tgggaagaat 180
 ctctcctttc tcctgcttga aggacagaca tgaattcaga ttcccggagg aggagtttga 240
 tggccaccag ttccagaaga ctcaagccat ctctgtcctc catgagatga tccagcagac 300
 cttcaatctc ttcagcacag aggactcatc tgctgcttgg gaacagagcc tcctagaaaa 360
 attttccact gaactttacc agcaactgaa tgacctggaa gcatgtgtga tacaggaggt 420
 tgggggtggaa gagactcccc tgatgaatga ggacttcac ctggctgtga ggaaatactt 480
 ccaaagaatc actctttatc taacagagaa gaaatacagc ccttgtgcct gggagggtgt 540
 cagagcagaa atcatgagat ccttctcttt ttcaacaaac ttgaaaaaag gattaaggag 600
 gaaggattga aaactgggtc atcatggaaa tgattctcat tgactaatgc atcatctcac 660
 actttcatga gttcttccat ttcaaagact cacttctata accaccacaa gttgaatcaa 720
 aattttccaaa tgttttc 737

<210> 24
 <211> 1933
 <212> DNA
 <213> Homo sapiens

<400> 24
 gatctggtaa acctgaagca aatatagaaa cctatagggc ctgacttcct acataaagta 60
 aggagggtaa aaatggaggc tagaataagg gttaaaattt tgcttctaga acagagaaaa 120
 tgattttttt catatatata tgaatatata ttatatatac acatatatac atatattcac 180
 tatagtgtgt atacataaat atataatata tatattgtta gtgtagtgtg tgtctgatta 240
 ttacatgca tatagtatat acacttatga ctttagtacc cagacgtttt tcatttgatt 300
 aagcattcat ttgtattgac acagctgaag tttactggag tttagctgaa gtctaatagca 360
 aaattaatag attgttgtca tcctcttaag gtcataaggga gaacacacaa atgaaaacag 420
 taaaagaaac tgaaagtaca gagaaatgtt cagaaaatga aaaccatgtg tttcctatta 480
 aaagccatgc atacaagcaa tgtcttcaga aaacctaggg tccaaggta agccatatcc 540
 cagctcagta aagccaggag catcctcatt tcccaatggc cctcctgttc cctctactgg 600

cagccctagt gatgaccagc tatagccctg ttggatctct gggctgtgat ctgcctcaga 660
 accatggcct acttagcagg aacaccttgg tgcttctgca ccaaagagg agaattctccc 720
 ctttcttggtg tctcaaggac agaagagact tcaggttccc ccaggagatg gtaaaaggga 780
 gccagttgca gaaggcccat gtcattgtctg tcctccatga gatgctgcag cagatcttca 840
 gcctcttcca cacagagcgc tcctctgctg cctggaacat gaccctccta gaccaactcc 900
 aactgggact tcattcagcaa ctgcaacacc tggagacctg cttgctgcag gtagtgggag 960
 aaggagaatc tgctggggca attagcagcc ctgcactgac cttgaggagg tacttccagg 1020
 gaatccgtgt ctacctgaaa gagaagaaat acagcgactg tgcttgggaa gttgtcagaa 1080
 tggaaatcat gaaatccttg ttcttatcaa caaacatgca agaaagactg agaagtaaag 1140
 atagagacct gggctcatct tgaaatgatt ctcatgtatt aatttgccat ataacacttg 1200
 cacatgtgac tctggtcaat tcaaaagact cttatttcgg ctttaatcac agaattgact 1260
 gaattagttc tgcaaatact ttgtcgggtat attaagccag tatatgttaa aaagacttag 1320
 gttcaggggc atcagtccct aagatgttat ttatttttac tcattttattt attcttacat 1380
 tttatcatat ttatactatt tatattctta tataacaaat gtttgccttt acattgtatt 1440
 aagataacaa aacatgttca gctttccatt tgggttaaata ttgtattttg ttatttatta 1500
 aattattttc aaacaaaact tcttgaagtt atttattcga aaacaaaat ccaaacacta 1560
 gttttctgaa ccaaatacag gaatggacgg taatatacac ttacctattc attcattcca 1620
 tttacataat atgtataaag tgagtatcaa agtggcatat tttggaattg atgtcaagca 1680
 atgcaggtgt actcattgca tgactgtatc aaaatatctc atgtaaccaa taaatatata 1740
 cacttactat gtatcccaca aaaattaaaa agttattttta aaaaagaaat acaggtgaat 1800
 aaacacagtt tctttccgtg ttgaagagct ttcatcttta caggaaaaga aacagtaaag 1860
 atgtaccaat ttcgcttata tgaaacacta caaagataag taaaagaaaa tgatgttctc 1920
 atactagaag ctt 1933

<210> 25
 <211> 734
 <212> DNA
 <213> Homo sapiens

<400> 25
 tgggtgacag cctcagagtg tttcttctgc tgacaaagac cagagatcag gaatgaaact 60
 agacatgact ggggactgca cgccagtgtt ggtgctgatg gccgcagtgc tgaccgtgac 120
 tggagcagtt cctgtcgcca ggctccacgg ggctctcccc gatgcaaggg gctgccacat 180

```

agcccagttc aagtccctgt ctccacagga gctgcaggcc ttttaagaggg ccaaagatgc 240
cttagaagag tcgcttctgc tgaaggactg caggtgccac tcccgcctct tccccaggac 300
ctgggacctg aggcagctgc aggtgaggga ggcgcccatg gctttggagg ctgagctggc 360
cctgacgctg aaggttcttg aggccaccgc tgacactgac ccagccctgg tggacgtctt 420
ggaccagccc cttcacaccc tgcaccatat cctctcccag ttccgggcct gtatccagcc 480
tcagcccacg gcagggccca ggaccgggg cgcctccac cattggctgt accggtcca 540
ggaggcccca aaaaaggagt cccctggctg cctcgaggcc tctgtcacct tcaacctctt 600
ccgcctcctc acgcgagacc tgaattgtgt tgccagtggg gacctgtgtg tctgacctc 660
ccaccagtca tgcaacctga gattttatth ataaattagc cacttgtctt aattttattgc 720
caccagtcg ctat 734

```

```

<210> 26
<211> 594
<212> DNA
<213> Homo sapiens

```

```

<400> 26
gacatgaccg gggactgcat gccagtgtg gtgctgatgg ccgcagtgt gaccgtgact 60
ggagcagttc ctgtcgccag gctccgcggg gctctcccgg atgcaagggg ctgccacata 120
gcccagttca agtccctgtc tccacaggag ctgcaggcct ttaagagggc caaagatgcc 180
ttagaagagt cgcttctgct gaaggactgc aagtgccgct cccgcctctt ccccaggacc 240
tgggacctga ggcagctgca ggtgaggagg cgccccgtgg ctttggaggc tgagctggcc 300
ctgacgctga aggttctgga ggccaccgct gacactgacc cagccctggg ggatgtcttg 360
gaccagcccc ttcacaccct gcaccatata ctctcccagc tccgggcctg tatccagcct 420
cagcccacgg cagggcccag gaccgggggc cgcctccacc attggctgca ccggctccag 480
gaggcccca aaaaaggagt cccctggctg ctcgaggcct ctgtcacctt caacctcttc 540
cgcctcctca cgcgagacct gaattgtgtt gccagcgggg acctgtgtgt ctga 594

```

```

<210> 27
<211> 594
<212> DNA
<213> Homo sapiens

```

```

<400> 27
gacatgaccg gggactgcat gccagtgtg gtgctgatgg ccgcagtgt gaccgtgact 60
ggagcagttc ctgtcgccag gctccgcggg gctctcccgg atgcaagggg ctgccacata 120

```

58

```

gccagttca agtcctgtc tccacaggag ctgcaggcct ttaagagggc caaagatgcc      180
ttagaagagt cgcttctgct gaaggactgc aagtgccgct cccgcctctt cccaggacc      240
tgggacctga ggcagctgca ggtgagggag cgccccgtgg ctttgagggc tgagctggcc      300
ctgacgctga aggttctgga ggccaccgct gacactgacc cagccctggg ggatgtcttg      360
gaccagcccc ttcacacctt gcaccatata ctctcccaga tccgggcctg tatccagcct      420
cagcccacgg cagggcccag gaccgggggc cgctccacc attggctgca ccggctccag      480
gaggcccca aaaaggagtc ccttggtgct ctcgaggcct ctgtcacctt caacctcttc      540
cgctcctca cgcgagacct gaattgtgtt gccagcgggg acctgtgtgt ctga          594

```

```

<210> 28
<211> 856
<212> DNA
<213> Homo sapiens

```

```

<400> 28
aattaccttt tcactttaca cacatcatct tggattgccc attttgcggtg gctaaaaagc      60
agagccatgc cgctggggaa gcagttgcga tttagccatg gctgcagctt ggaccgtggt      120
gctggtgact ttggtgctag gcttggccgt ggcaggccct gtccccactt ccaagcccac      180
cacaactggg aagggtgctc acattggcag gttcaaactc ctgtcaccac aggagctagc      240
gagcttcaag aaggccaggg acgccttgga agagtcactc aagctgaaaa actggagttg      300
cagctctcct gtcttccccg ggaattggga cctgaggctt ctccagggtga gggagcgccc      360
tgtggccttg gaggtgagc tggccctgac gctgaaggtc ctggaggccg ctgctggccc      420
agccctggag gacgtcctag accagcccct tcacaccctg caccacatcc tctcccagct      480
ccaggcctgt atccagcctc agcccacagc agggcccagg ccccgggggc gcctccacca      540
ctggctgcac cggtccagg agggcccca aaaggagtcc gctggctgcc tggaggcatc      600
tgtcaccttc aacctcttcc gcctcctcac gcgagacctc aaatatgtgg ccgatgggaa      660
cctgtgtctg agaacgtcaa cccacctga gtccacctga cccccacac cttatttatg      720
cgctgagccc tactccttcc ttaatttatt tcctctcacc cttatttat gaagctgcag      780
ccctgactga gacatagggc tgagtttatt gttttacttt tatacattat gcacaaataa      840
acaacaagga attgga          856

```

```

<210> 29
<211> 607
<212> DNA
<213> Artificial Sequence

```

<220>

<223> Synthetic DNA encoding human interferon-like peptide

<400> 29

```

aattcaccgt gcgtgttgac tattttacct ctggcgggtga taatgaaatc aattgtgagc      60
ggataacaat taaaaattaa ggaggatcac tatctgtgat ttgccgcaga ctcatagctt      120
cggcaaccga acagcactca tcctcttggc ccaattggcc caaatgggtc gcatttcctt      180
gttctcgtgc cttaaagacc gccacgattt cgggtttcca caggaagagt ttgacggcaa      240
tcagttccaa aaggctcagg caatctcggg tctgcatgag atgattcagc aaatcttcaa      300
cctgtttctc actaaagact ctteggctgc ttggaacgaa tccttgcttg ataaattctc      360
cactgaactc tatcagcaac tgaacgtact tgaagcttgc gttatccagg aggtaggcgt      420
tgaagagact ccgcttatga atgttgagtc taccctggct gtatctaagt attttcaccg      480
aattaccctc tacctcagtg agaagaaata ttcaccgtgt gcgtgggaaa ttgtgagagc      540
cgaaatcatg cgttccttga ctcttctgac caacctccag gaacgcctgc gtaataaaga      600
ctaatag                                           607

```

<210> 30

<211> 611

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic DNA encoding human interferon-like peptide

<400> 30

```

aattcaccgt gcgtgttgac tattttacct ctggcgggtga taatgaaatc aattgtgagc      60
ggataacaat tattacccaa cttgaggaat ttataatgtg tgatttacca caaacccata      120
gcttgggcaa ccgaagagca ctcatcctgt tggcccaatt ggcccaaatt ggctcgcatTT      180
ccctgttctc gtgccttaaa gaccgccacg atttcgggtt tccacaggaa gagtttgacg      240
gcaatcagtt ccaaaaggct caggcaatct cggttctgca tgagatgatt cagcaaattct      300
tcaacctgtt ctccactaaa gactgttcgg ctgcttgga cgaatccttg cttgataaat      360
tctccactga actctatcag caactgaacg tacttgaagc ttgcgttata caggaggtag      420
gcgttgaaga gactccgctt atgaatgttg agtctatcct ggctgtatct aagtattttc      480
accgaattac cctctacctc agtgagaaga aatattcacc gtgtgcgtgg gaaattgtga      540
gagccgaaat ctagcgttcc ctgactcttc tgacgaacct ccaggaacgc ctgcgtaata      600
aagactaata g                                           611

```

60

<210> 31
 <211> 2035
 <212> DNA
 <213> Homo sapiens

<400> 31
 cgagccccgc cgaaccgagg ccacccggag ccgtgcccag tccacgccgg ccgtgcccgg 60
 cggccttaag aaccaggcaa cctctgcctt cttccctctt ccactcggag tcgcgctccg 120
 cgcgccctca ctgcagcccc tgcgtgcgcg ggaccctcgc gcgcgaccag ccgaatcgct 180
 cctgcagcag agccaacatg cccatcactc ggatgcgcat gagaccctgg ctagagatgc 240
 agattaattc caaccaaadc cgggggctca tctggattaa taaagaggag atgatcttcc 300
 agatcccatg gaagcatgct gccaaagcat gctgggacat caacaaggat gcctgtttgt 360
 tccggagctg ggccattcac acaggccgat acaaagcagg ggaaaaggag ccagatccca 420
 agacgtggaa ggccaacttt cgctgtgcc a tgaactccct gccagatata gaggaggtga 480
 aagaccagag caggaacaag ggcagctcag ctgtgcgagt gtaccggatg cttccacctc 540
 tcaccaagaa ccagagaaaa gaaagaaagt cgaagtccag ccgagatgct aagagcaagg 600
 ccaagaggaa gtcattgtgg gattccagcc ctgatacctt ctctgatgga ctcagcagct 660
 ccactctgcc tgatgaccac agcagctaca cagttccagg ctacatgcag gacttggagg 720
 tggagcaggc cctgactcca gcactgtcgc catgtgctgt cagcagcact ctccccgact 780
 ggcacatccc agtggaaagt gtgccggaca gcaccagtga tctgtacaac ttccagggtgt 840
 cccccatgcc ctccacctct gaagctacaa cagatgagga tgaggaaggg aaattacctg 900
 aggacatcat gaagctcttg gagcagtcgg agtggcagcc aacaaacgtg gatgggaagg 960
 ggtacctact caatgaacct ggagtccagc ccacctctgt ctatggagac tttagctgta 1020
 aggaggagcc agaaattgac agcccagggg gggatattgg gctgagtcta cagcgtgtct 1080
 tcacagatct gaagaacatg gatgccacct ggctggacag cctgctgacc ccagtccggt 1140
 tgccctccat ccaggccatt ccctgtgcac cgtagcaggg ccctggggcc cctcttattc 1200
 ctctaggcaa gcaggacctg gcatcatggt ggatatggtg cagagaagct ggacttctgt 1260
 gggccccctca acagccaagt gtgacccccc tgccaagtgg ggatgggcct ccctccttgg 1320
 gtcattgacc tctcagggcc tggcaggcca gtgtctgggt ttttcttgtg gtgtaaagct 1380
 ggccctgcct cctgggaaga tgaggttctg agaccagtgt atcaggtcag ggacttggac 1440
 aggagtcagt gtctggcttt ttctcttgag ccagctgcc tggagagggt ctgcgtgtca 1500
 ctggctggct cctaggggaa cagaccagtg accccagaaa agcataaac caatcccagg 1560
 gctggctctg cactaagcga aaattgcact aaatgaatct cgttccaaag aactaccct 1620

tttcagctga gccctgggga ctgttcctaaa gccagtgaat gtgaaggaaa ctccccctcct	1680
tcgggggcaat gctccctcag cctcagagga gctctaccct gctccctgct ttggctgagg	1740
ggcttgggaa aaaaacttgg cacttttttcg tgtggatcct gccacatttc tgatcagagg	1800
tgtacactaa catttcccc gagctcttgg cctttgcatt tatttataca gtgccttgct	1860
cggggcccac caccctca agccccagca gccctcaaca ggcccaggga gggaagtgtg	1920
agcgccttgg tatgacttaa aattggaaat gtcactaac cattaagtca tgtgtgaaca	1980
cataaggacg tgtgtaaata tgtacatttg tctttttata aaaagtaaaa ttggtt	2035

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2005/014626

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K38/21 A61K9/00 A61K9/127 A61K9/51 A61P35/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, EMBASE, MEDLINE, BIOSIS, WPI Data, PAJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 2004/074314 A (UNIVERSITY OF SOUTH FLORIDA; MOHAPATRA, SHYAM, S) 2 September 2004 (2004-09-02) claims 1-4	23-25
Y	WO 03/092618 A (UNIVERSITY OF SOUTH FLORIDA; MOHAPATRA, SHYAM, S; BEHERA, ARUNA, K) 13 November 2003 (2003-11-13) claims 1-54 <div style="text-align: center; margin-top: 10px;">----- -/--</div>	26-31
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
° Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
28 September 2005	12/10/2005	
Name and mailing address of the ISA	Authorized officer	
European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Ganschow, S	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2005/014626

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KUMAR M ET AL: "ROLE OF MUCOSAL IFN-GAMMA GENE TRANSFER ALLERGIC SENSITIZATION AND RSV INFECTION" JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, MOSBY - YEARLY BOOK, INC, US, vol. 109, no. 1, January 2002 (2002-01), page S43, XP008033852 ISSN: 0091-6749	26-31
Y	abstract	1-22
Y	----- LENGYEL P: "Tumor-suppressor genes: News about the interferon connection" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA 1993 UNITED STATES, vol. 90, no. 13, 1993, pages 5893-5895, XP002347048 ISSN: 0027-8424 the whole document	1-22
A	----- WO 01/22970 A (THE CLEVELAND CLINIC FOUNDATION; SILVERMAN, ROBERT, H; RUSCH, LORRAINE) 5 April 2001 (2001-04-05) claims 1,2 -----	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2005/014626

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-22 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2005/014626

Patent document cited in search report			Publication date	Patent family member(s)	Publication date
WO 2004074314	A		02-09-2004	NONE	
WO 03092618	A		13-11-2003	AU 2003225281 A1	17-11-2003
				CA 2484251 A1	13-11-2003
				EP 1579001 A2	28-09-2005
WO 0122970	A		05-04-2001	AU 1494801 A	30-04-2001
				CA 2388025 A1	05-04-2001
				EP 1229920 A1	14-08-2002
				JP 2003510281 T	18-03-2003